



The microbiome: An emerging key player in aging and longevity

Minhoo Kim ^a, Bérénice A. Benayoun ^{a, b, c, d, *}

^a Leonard Davis School of Gerontology, University of Southern California, Los Angeles, CA, 90089, USA

^b USC Norris Comprehensive Cancer Center, Epigenetics and Gene Regulation, Los Angeles, CA, 90089, USA

^c Molecular and Computational Biology Department, USC Dornsife College of Letters, Arts and Sciences, Los Angeles, CA, 90089, USA

^d USC Stem Cell Initiative, Los Angeles, CA, 90089, USA

ARTICLE INFO

Article history:

Received 1 June 2020

Received in revised form

15 July 2020

Accepted 17 July 2020

Available online 21 July 2020

Keywords:

Microbiome

Aging

Sex-dimorphism

ABSTRACT

Revolutionary advancements of high-throughput sequencing and metagenomic tools have provided new insights to microbiome function, including a bidirectional relationship between the microbiome and host aging. The intestinal tract is the largest surface in the human body that directly interacts with foreign antigens – it is covered with extremely complex and diverse community of microorganisms, known as the gut microbiome. In a healthy gut, microbial communities maintain a homeostatic metabolism and reside within the host in a state of immune tolerance. Abnormal shifts in the gut microbiome, however, have been implicated in the pathogenesis of age-related chronic diseases, including obesity, cardiovascular diseases and neurodegenerative diseases. The gut microbiome is emerging as a key factor in the aging process. In this review, we describe studies of humans and model organisms that suggest a direct causal role of the gut microbiome on host aging. Additionally, we discuss sex-dimorphism in the gut microbiome and its possible roles in age-related sex-dimorphic phenotypes. We also provide an overview of widely used microbiome analysis methods and tools which could be used to explore the impact of microbiome remodeling on aging.

© 2020 The Authors. Production and hosting by Elsevier B.V. on behalf of KeAi Communication Co., Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Over a century ago, Elie Metchnikoff proposed that age-related dysfunction could result from increased colon permeability-driven chronic system inflammation [1]. Recent advances in DNA sequencing technologies have allowed investigation of the composition and functional dynamics of complex microbial communities with great resolution and without the need for cultivation [2]. During the past two decades, microbiome research thrived to establish a causal relationship between the microbiome and host aging (Reviewed in Ref. [3–6]).

The microbiota consists of all the microbes (*i.e.* bacteria, archaea, viruses, protozoa and fungi) and a distinct profile of microbiota is found on all host surfaces that are in direct contact with the outside environment (*e.g.* gut, skin, mouth, vagina, *etc.*) [7–15]. The microbiota performs essential functions that contribute to the physiology of the host through a symbiotic relationship [16,17].

Consequently, perturbations of microbiomes have been proposed to exert negative effects on the host organism. The gut microbiome remains the most extensively studied microbiome and will be the main focus of our review.

Intestinal mucosa is the largest surface of the body that directly interacts with environmental antigens. Thus, the intestinal mucosal immune system monitors the gut environment through a variety of pattern-recognition receptors and is in active communication with the systemic immune system via the local mesenteric lymph nodes [18,19]. The adult human gut microbiome, composed of approximately 10^{13} to 10^{14} micro-organisms, plays various essential roles in the host including degradation of food, lipid storage and metabolism, vitamin synthesis, suppression of harmful microbial species and maintenance of intestinal barrier integrity [20–25]. Dysbiosis of the gut microbiome is associated with defects in gut barrier integrity and enhanced pro-inflammatory cytokines [26,27]. Thus, aberrant alterations of the gut microbiome have been attributed to pathogenesis of various metabolic diseases including adiposity, insulin resistance, atherosclerosis and cardiovascular disease, as well as multiple sclerosis, depression and anxiety [28–32].

* Corresponding author. Leonard Davis School of Gerontology, University of Southern California, Los Angeles, CA, 90089, USA.

E-mail address: berenice.benayoun@usc.edu (B.A. Benayoun).

In this review, we describe changes to the gut microbiome throughout lifespan of the host and key findings that implicate a central role of the gut microbiome in host aging. In addition, we discuss microbiome-relevant biological factors (e.g. sex) that may contribute to aging. Finally, we provide an overview of microbiome collection considerations, data analysis pipelines and potential confounding factors, that must be considered when analyzing and interpreting microbiome data.

2. The microbiome in response to aging and pro-longevity interventions

2.1. The aging gut microbiome: the human side

The microbiota co-evolves with its host and thus the composition of the microbial community within the intestinal tract fluctuates throughout lifespan, in response to genetic and environmental stimuli [3,33–35]. Based on recent studies that reported the presence of bacteria in the placenta, amniotic cavity and umbilical cord, microbial colonization may initiate as early as *in utero* [36–38]. During infancy, the gut microbiome undergoes significant fluctuations, which is namely driven by factors including delivery method, feeding, antibiotic exposure, maternal diet and environmental factors [39–41]. Colonization of microbial species in the gastrointestinal tract during early stages of life is reported to affect later health of the host organism [42]. Nonetheless, the microbiome composition reaches a stable structure after the first three years, its profile resembling that of an “adult-like” microbiome [43–45]. After stable recolonization of the microbiome, diet becomes a major force shaping the microbiome composition of the host throughout early adulthood [46,47].

In general, healthy adults are reported to present with high levels of bacteria from the *Bacteroidetes* and *Firmicutes* phyla, and relatively lower proportions of the *Proteobacteria*, *Actinobacteria*, *Fusobacteria* and *Verrucomicrobia* phyla [8,48–50]. *Bacteroidetes* found in the gut mainly functions in polysaccharide metabolism and calorie absorption, whereas *Firmicutes* are important for production of Short-Chain Fatty Acids (SCFAs) [51–55]. The *Firmicutes/Bacteroidetes* (F/B) ratio is reported to increase from birth to adulthood and studies show that high F/B ratios are associated to a dysbiotic microbiome [34,56]. Studies have shown that the F/B ratio can be used as an important indicator of gut microbiome state and thus host health [57–60].

Clinical studies have reported significant differences in microbial composition between young and elderly human subjects [6,61]. A key transition from healthy adult to elderly microbiota is characterized by a decrease in microbial diversity. Reduced microbiota diversity in aged individuals have been suggested to result in the expansion of distinct groups of bacteria which has been implicated on the development of age-associated type 1 diabetes mellitus, rheumatoid arthritis and colitis [62–64]. However, whether reduced microbiota diversity directly impacts host aging, or is a mere bystander, remains poorly understood.

Generally, in aged individuals, a decrease in *Bifidobacterium* and *Lactobacillus*, and increase in *Enterobacteriaceae* are observed [39,65,66]. Such changes in the microbiome structure are believed to result from changed lifestyle, dietary pattern, reduced mobility, weakened immune strength, reduced intestinal functionality, changes in gut morphology, use of medication, recurrent infections and more [27,34,35,39,61,65,67]. However, it is important to note that these generalizations do not apply to certain aged groups from different geological locations or genetic backgrounds [67,68]. Interestingly, in centenarians and supercentenarians, health-associated bacteria genera, including *Bifidobacteria* and *Christensenella*, are especially abundant [69,70]. Although these

observations are correlative, studies in model organisms support pro-longevity and pro-health effects of these microbes [71–73]. For example, supplementation of *Bifidobacteria* to *C. elegans* resulted in reduced accumulation of lipofuscin, a marker of aging, improved locomotor function and increased longevity [72]. Additionally, transplantation of *Christensenella* to germ-free mice has been shown to amend obese-associated microbiome and reduce weight gain [73].

2.2. The aging microbiome in model organisms

Baseline microbial composition of the gut microbiota varies across species and taxa [26,74–76]. However, similar to what has been observed in humans, extensive remodeling of the gut microbiome during aging has also been observed in a number of model organisms, spanning *Drosophila melanogaster*, the African turquoise killifish *Nothobranchius furzeri* and mice [26,74–77].

In *D. melanogaster*, the aging gut microbiome is characterized by an expansion of *Gammaproteobacteria* [74,75], and microbiota transplantation from aged donors to young flies leads to reduced longevity. Metagenomics analysis also showed that age-related changes in *Drosophila* microbial species were somewhat similar to that observed in human inflammatory disorder patients and aged human gastrointestinal tract [75]. For example, increased levels of *Enterobacteriaceae*, the most abundant family of *Gammaproteobacteria*, were also observed in aged humans and mouse model of colitis [78,79]. In a study of the aging microbiome in the African turquoise killifish, the microbiota of young individuals was found to be more enriched in species from the *Bacteroidetes*, *Firmicutes*, and *Actinobacteria* phyla, whereas the aged microbiota was enriched for species from the *Proteobacteria* phylum [76]. Interestingly, similar to observations in humans, the aging killifish microbiome was also characterized by decreased diversity of the gut microbial community [76]. Additionally, transplantation of microbiome from young to middle-aged killifish improved locomotion and longevity of recipient subjects [76]. The mouse aging microbiome showed a number of shifts in relative abundance of bacteria phyla, including increased presence of *Clostridium* and decreased levels of *Lactobacillaceae* as observed with human aging [80]. Increased abundance of *Clostridium* was also observed in the aging gut microbiome of rats, although (contrary to humans) rats seemed to acquire increased microbial diversity throughout life [81]. Interestingly, the relative proportion of *Firmicutes* and *Bacteroidetes* is also altered with aging in the gut microbiome of aging mice [26]. A direct role of the microbiome in promoting overall health in mammals is also suggested by the fact that fecal microbiota transplants from wild-type mice can significantly improve the health and lifespan of progeroid mice [82]. Together, these studies provide a strong rationale for microbiome-based interventions against age-related decline and pathologies.

2.3. Effects of pro-longevity interventions on the aging microbiome

Modulation of the microbiota is emerging as a potential mechanism underlying pro-health and longevity effects of various interventions (Table 1). Interestingly, a number of pro-longevity interventions seem to have rejuvenating effects on the microbiome. A recurrent effect is the expansion of bacteria from the *Lactobacillaceae* taxa, which occurs in the context of independent interventions [83,84]. Interestingly, a study reported that weight loss in the context of calorie restriction in mice seems to require an intact microbiome [83]. Thus, it will be important to determine whether microbial community remodeling in the context of pro-longevity interventions is a mere bystander, or an actual mediator of pro-health effects.

Table 1

Impact of pro-longevity interventions on the aging microbiome.

Pro-longevity intervention	Organism	Site	Effect on microbiome	Microbiome profiling	Reference
Calorie restriction	<i>M. musculus</i>	Gut	- Gut microbiota required for CR-induced weight loss - Significant increases in <i>Lactobacillus</i> , <i>Bifidobacterium</i> - Decreased B/F ratio	V4 16S V3–V4 16S	[83,84,219]
	<i>R. norvegicus</i>	Gut	- Increase of <i>Lactobacillus</i> - Increased B/F ratio - Changes in microbial SCFA production (Increased propionogenesis, decreased butyrogenesis and acetogenesis)	V4 16S Full length 16S	[220,221]
Dwarfism (Ames)	<i>M. musculus</i>	Gut	- Increased B/F ratio	V4 16S	[219]
	<i>C. elegans</i>	Gut (food source)	- Changes the <i>E. coli</i> metabolism of folate and methionine - <i>E. coli</i> required for longevity extension	N/A	[222]
Metformin	<i>H. sapiens</i>	Gut	- Increased <i>E. coli</i> abundance - Increased production of SCFAs	V4 16S Metagenomics	[223–225]
	<i>M. musculus</i>	Gut	- Increased B/F ratio - Increased abundance of <i>Lactobacillus</i> - Rejuvenation of the oral microbiome	V4 16S	[226]
Rapamycin	<i>M. musculus</i>	Oral	- Increased prevalence of segmented filamentous bacteria	V4 16S	[13]
	<i>M. musculus</i>	Gut	- Remodeling of specific OTUs - No change in B/F ratio - Renders microbiome more similar to that of HFD-treated mice	V4 16S Full length 16S (Sequencing and PhyloChip)	[227–229]
Resveratrol	<i>M. musculus</i>	Gut	- No change in B/F ratio - Reverses HFD-induced changes in bacterial abundances	Full length 16S	[229]
Mediterranean diet	<i>H. sapiens</i>	Gut	- Increased taxa associated to lower frailty in aged humans	V3–V4 16S	[230]
Acarbose	<i>M. musculus</i>	Gut	- Predicted increase in microbial SCFA production - Increased abundance in <i>Muribaculaceae</i> - Increase in microbial SCFAs, including propionate	V4 16S	[231]

CR: Calorie restriction.

B/F: Bacteroidetes/Firmicutes.

SCFA: Short-chain fatty acid.

OTU: Operational taxonomic unit.

HFD: High-fat diet.

3. A bidirectional relationship between the gut microbiome and aging?

During the past two decades, studies have provided evidence that age-associated shifts in the gut microbiome contributes to increased predisposition of aged individuals to certain diseases, including cardiovascular diseases, cancer, obesity, cancers, diabetes and neurodegenerative diseases [3,4,85–87]. Aging is a complicated process that affects physiological, metabolic and immunological functions of the organism and thus is accompanied by inflammation and metabolic dysfunctions [88]. The overall age-related increase in chronic inflammation and deterioration of systemic immune system led to coining the term “inflamm-aging” [89]. A direct causal role of the gut microbiome on host aging has been suggested by a number of studies using various experimental models [76,90]. In this section, we discuss studies that suggest the existence of a bidirectional relationship between the gut microbiome and host aging.

3.1. Interaction between the host immune system and the gut microbiome

Through millions of years of evolution, the host and its surrounding microbial environment have co-evolved into a complex organism [17,18,91]. Microbes beneficial to the host are able to reside within the host in a state of immune tolerance, whereas those that exert a pathogenic effect activates robust immune responses of the host [17]. The symbiotic co-existence between the host and microbiota is feasible due to the anatomical separation of microbial species from the host by a physical barrier. The intestinal barrier is responsible for adjusting metabolic homeostasis and systemic antimicrobial responses by detecting microbial-cell components and metabolites through its extensive repertoire of

innate immune receptors [92–96]. For example, activation of pattern-recognition receptors (e.g. Toll-like receptors) by the gut microbe or its products induces the production of antimicrobial peptides and mucus [92]. Perturbations of such receptors have been reported to result in intestinal inflammation and susceptibility to enteric infections [97].

Relevant to aging, decline of the immune system in the aged intestinal epithelium have been suggested to contribute to age-onset dysbiosis [98,99]. An important characteristic of age-onset dysbiosis is reduced microbiota diversity, which is suggested to lead to an expansion of distinct groups of bacteria [39,100,101]. Concurrently, bacteria that is reported to be involved in maintenance of immune tolerance in the gut, such as *Bifidobacteria* and *Lactobacilli*, are found in reduced level in aged groups, whereas those that are found in increased levels, such as *Enterobacteriaceae* and *Clostridium*, are involved in infection and intestinal inflammation stimulation [27,66,102,103]. Together, these studies suggest that the host immune system shapes not only the host's immune response to microbiome changes, but also the structure of the microbiome itself [104].

Cumulative evidence has implicated a close functional relationship between the immune system of the host and the microbiome, to an extent that the gut microbiome is important for proper development and expansion of intestinal mucosal and systemic immune system [105,106]. Supporting the notion that the microbiome can directly shape the immune states of the host, the transcriptomic profile of African turquoise killifish guts derived from animals that received young or old gut microbiota transplants showed clear differences, especially in expression of immune-related genes [76]. Interestingly, studies using germ-free mice models also suggest a bidirectional relationship between the host immune system and the gut microbiome. Germ-free mice showed significant alterations in innate immune system composition

compared to classical Specific-Pathogen Free [SPF] mice, including deficiencies in macrophage, monocyte and neutrophil populations [107]. Such alterations of the immune system in germ-free mice were partially rescued when mice were treated with specific bacteria and/or bacterial components (*i.e.* bacterial polysaccharide), demonstrating a direct role of the gut microbiome on immune system establishment of the host [108]. Interestingly, experiments using germ-free mice also revealed that, in addition to regulating the abundance of immune cells, the microbiome may also regulate bactericide properties of macrophages [26].

3.2. Increased intestinal barrier permeability with age

Increased permeability of the intestinal barrier with age has been described across animal species, including worms, flies, mice and rats [26,74,75,109–111]. Age-related deterioration of intestinal barrier function has been proposed to result in leakage of gut microbes into the systemic circulation, and ultimately lead to increased antigenic load and systemic immune activation [112,113] (Fig. 1). For example, age-associated remodeling of the gut

microbiome in mice was shown to result in increased production of pro-inflammatory cytokines and intestinal barrier failure [26]. Consistently, the blood of aged mice contained increased levels of muramyl-dipeptide, a component of Gram-positive and Gram-negative bacteria cell wall [26]. Additionally, in a clinical study on aged type 2 diabetes patients, live gut bacteria were found to translocate into the blood stream, suggesting perturbations of the intestinal barrier integrity [114]. In *Drosophila*, the age-related increase in *Gammaproteobacteria* was suggested to lead to increased intestinal permeability, inflammation and mortality [74,75]. The study showed that regardless of chronological age, intestinal dysbiosis serves as an indicator of age-onset mortality in flies [75].

A number of molecular mechanisms has been suggested to underlie intestinal barrier permeability with age. Mouse studies suggest that host cytokine signaling may play a key role in barrier function breakdown [26]. Indeed, TNF- α signaling was found to play a role in age-related intestinal barrier breakdown, as (i) *Tnfa* knock-out mice did not accumulate bacteria byproducts in their blood with aging, and (ii) anti- TNF- α therapy led to significant remodeling of the gut microbiota [26]. Mechanistically, age-




Fig. 1. The bidirectional relationship between the gut microbiome and aging

(Left panel) In a healthy gut, balanced microbial composition and intestinal barrier integrity maintains gut homeostasis and contains the microbiota in the intestinal lumen. Microbiota-derived metabolites, including SCFAs, participate in a feedback mechanism with the host immune system to fortify the barrier function, produce mucus and promote intestinal stem cell proliferation. An efficient immune system tolerates the host immune responses to avoid excessive activation. (Right panel) In gut dysbiosis (such as with aging), declined intestinal barrier integrity results in translocation of microbes and microbial particles through the intestinal epithelial cell lining. Reduced microbiota diversity leads to overgrowth of distinct microbes and metabolism instability. Aberrant levels of microbiota-derived metabolites instigate abnormal immune responses resulting in chronic inflammation. SCFA: Short-chain fatty acid.

associated epithelial tight-junctions permeability and declined function of Paneth cells of the intestinal mucosa have been speculated to result in intestinal barrier permeability [115,116]. However, further research will be needed to fully understand the mechanisms of age-associated increase in gut permeability.

3.3. Changes in production of microbiome-derived metabolites with age

The gut microbiome plays various essential roles in the host including degradation of food, lipid storage and metabolism, vitamin synthesis, suppression of harmful microbial species and maintenance of intestinal barrier integrity [25]. Microbiome-derived SCFAs, including butyrate, propionate, acetate and valerate, are important energy source for the epithelium and ultimately affects hypoxia-inducible factor-mediated fortification of the epithelial barrier [117]. Interestingly, a decline in SCFA levels, including that of butyrate, were observed in aged humans, whereas centenarians presented with a rearrangement in the population of specific butyrate-producing bacteria [70,118]. Additionally, the blood and intestine of germ-free mice presented with significantly lower levels of SCFAs compared to conventionally raised mice, supporting a role of the microbiota in regulating host SCFA levels [119–121]. For example, studies have shown that administration of butyrate restores the observed abnormal absorptive colonic motor activity and blood-brain barrier permeability in germ-free mice [122,123]. Microbiota-derived metabolites has also been reported to play a role in intestinal epithelial stem cell proliferation [4]. For example, butyrate and nicotinic acid, both by-products of the gut microbiota, are involved in suppression and promotion of stem cell proliferation in the colon, respectively [124]. In addition, microbiota-derived neurostimulators, including serotonin, glutamate, gamma-aminobutyric acid, have been reported to regulate proliferation of intestinal epithelial stem cells through the enteric nervous system [125]. Collectively, functional products of the microbiota have been implicated in the regulation of intestinal barrier integrity and function.

Other microbiota-derived metabolites have been shown to directly affect numerous systems of the host, although their functions in relation to host aging is in need of further investigation [126]. For example, TriMethylAmine n-Oxide [TMAO], a byproduct of microbial metabolism, is associated with cardiometabolic diseases, such as atherosclerosis and type 2 diabetes [127,128]. Interestingly, microbial metabolites may also affect the host through epigenetic alterations: indeed, microbiota-derived butyrate can affect the immune response of colonic macrophages through the inhibition of histone deacetylases [129,130]. The microbiome has also been shown to contribute to various neurological conditions through the so-called “microbiota-gut-brain axis” [131]. It is noteworthy that the microbiome can influence behavioral aspects of the host – the level of SCFAs is reported to affect feeding behavior of the host and thus energy homeostasis [132]. Given that diet is a key factor in remodeling of the microbiome structure, an integrative assessment of various physiological conditions of the host is required to precisely understand the functions and effects of the microbiome.

4. Sex-dimorphism in the gut microbiome, and possible impact on aging

Although aging is a conserved process across species and biological sex, accumulating evidence has shown that many age-related phenotypes are sex-dimorphic, and may thus modify aspects of aging between animals of opposite sex [133]. Concurrently, disparities between the sexes are observed in manifestation of

certain age-associated diseases, including obesity, multiple sclerosis and Alzheimer's disease [134,135]. However, due to experimental pragmatism, still few studies systematically evaluate how sex interacts with aging phenotypes, including age-related microbial dysbiosis. Fundamentally, key phenotypic sex differences are driven by genetic and/or hormonal mechanisms of the host [136]. Intriguingly, recent studies suggest that there may be substantial involvement of the gut microbiome in tuning sex-dimorphic phenotypes.

4.1. Sex-dimorphism in the gut microbiome

Mouse model studies have shown that the composition of the microbiome starts to diverge between male and female individuals after the onset of puberty [137,138]. As described above, the gut microbial composition of healthy human adults is reported to consist of high levels of *Bacteroidetes* and *Firmicutes* [8,48,49]. Interestingly, studies have shown females present with higher F/B ratio compared to that of males [16,139]. Additionally, *Proteobacteria*, *Veillonella* and *Blautia* are found in higher levels in females compared to males, but in much lower proportions [16,140]. Although increased F/B ratio is associated with gut dysbiosis, a systematic analysis of such disparities between the two sexes and understanding of its physiological implications are still lacking [34,56].

Interesting sexually dimorphic phenotypes have been described in studies using germ-free mice models. For example, Non-Obsese Diabetes [NOD] model female mice are more prone to spontaneously develop type 1 diabetes compared to NOD model male mice [137]. However, such difference between sexes disappeared when mice were raised in germ-free conditions [141]. In support of this finding, microbiota transplantation of conventionally raised NOD male mice microbiota to germ-free NOD female mice reduced the rate of type 1 diabetes incidence in the recipient mice [137].

4.2. Sex-dimorphism in host-microbiome communication?

In addition to observed sex differences in microbial communities' composition, emerging evidence is suggesting that the microbiome may potentiate the expression of sex-dimorphic phenotypes in the hosts. For instance, a recent study utilizing germ-free mice suggested that the microbiome is required to establish sex-dimorphic gene expression patterns in the liver [142]. Another study, also comparing germ-free to SPF mice, found that presence of microbes was required for sex-dimorphic regulation of lipid metabolism in the small intestine of mice [143]. Consistent with sex-dimorphic modulation of the immune system by the microbiome, the transcriptional response of adult microglia – the resident macrophages of the brain – to chronic (*i.e.* germ-free vs. SPF husbandry) or acute (*i.e.* antibiotic treatment) microbiota depletion was found to be sex-dimorphic [144]. Intriguingly, a recent study showed that microbiota depletion through antibiotic treatment rescued a number of brain phenotypes only in males in a mouse model of Alzheimer's disease [145]. Reestablishment of the microbiota reversed the rescue, supporting a direct implication of the microbiota in this phenomenon [145]. Thus, host responses to commensal microbes can be sex-dimorphic, revealing that the microbiome interacts with the biological sex of the host. However, how these sex-dimorphic interactions are modulated during aging remains largely unknown. Future studies investigating the impact of the microbiome on the aging process should systematically include sex as a variable to address this complex question.

4.3. Interactions between the microbiome and sex-steroid metabolism

The gut microbiome has been proposed to drive estrogen metabolism and regulate the proportions of recirculated and excreted estrogens and estrogen metabolites in the host organism [146–148]. The term “estrobolome” has been coined to define “the gene repertoire of the microbiota of the gut capable of metabolizing estrogens” [149,150]. Indeed, the human gut microbiome is able to hydrolyze estrogen sulfate and glucuronide conjugates [151]. Thus, through manipulation of the gut microbiome, circulating estrogen levels can be shifted in a dosage-dependent manner [148]. Consistently, in a recent study, germ-free female mice presented with significantly lowered levels of 17-β estradiol, the major form of estrogens in females, compared to conventionally raised mice [142]. In the same study, transcriptome analysis of sexual development marker genes and histological studies of follicle development in germ-free female mice indicated that sexual maturation is perturbed in microbiota-depleted mice [142].

Interestingly, estrogens have been shown to impact gut microbiome structure and contribute in gut homeostasis maintenance [152]. In a metabolic syndrome study, the microbiome structure of males and ovariectomized [OVX] females were observed to share similar profiles [153]. When the two test groups were supplemented with 17-β estradiol, both males and OVX females showed alteration of the gut microbiome and suppression of Western diet-induced obesity phenotypes. Collectively, these findings demonstrate a close bidirectional relationship between the gut microbiome and female sex hormones in affecting host health.

5. Microbiome data analysis: experimental and analytical “omics” pipelines

Traditionally, research on microbial interactions was focused on single pathogenic organisms through culture-based methods that capture only a small proportion of the bacterial microbiota [154]. However, recent findings suggest that disease pathogenesis is dependent not only on single pathogens, but also on global changes in the host microbiome [155,156]. Advancements in next-generation sequencing techniques have enabled culture-independent analyses to capture the global changes in the microbiome. In addition, the advent of various model organisms and experimental tools, including germ-free rodent models and microbiota transplantation methods, have helped characterize microbial communities as key factors in not only dietary metabolism and host nutrition, but also in the pathogenesis of a number of chronic age-associated disease, including diabetes, cardiovascular diseases and neurodegenerative disorders [157–161].

As the field expands, microbiome analysis methods and standards are rapidly advancing to allow accurate characterization and interpretation of the microbiome data. This section will provide a primer on microbial sample collection guidelines and widely used microbiome data analysis methods: marker gene, metagenomics and metatranscriptomics analysis (Table 2).

5.1. Guidelines for microbiome sample collection

For human studies, oral, skin and vaginal samples are generally collected by a physician during a clinic visit – microbial samples can be collected by swabbing the appropriate area using a sterile soft cotton tip or nylon swab [162]. For model organisms, the same type of swab can be used for sample collection. Samples should be immediately flash-frozen and stored at -80 °C until further processing [162]. Among various microbial samples, fecal sample collection for gut microbiome analysis presents with the most

challenges because on demand collection of fecal samples is difficult. For human fecal samples, various transportation kits, including the Fisherbrand™ Commode Specimen Collection System (Fisher Scientific), OMNIgene Gut kit (DNA Genotek) and Cary Blair Transport Medium (Remel), have been developed in order to preserve microbial composition during shipping from site of sample collection to laboratories for further analysis [163–165].

For mouse studies, gut microbiome samples can be collected by picking freshly defecated fecal pellets or extracting fecal pellets from the distal colon after euthanasia. When collecting from the distal colon, fecal pellets need to be homogenized in order to ensure even distribution of microbial species of the colon. Immediate freezing of fecal samples is crucial – storage of microbiome samples at room or higher temperature for extended times results in expansion of specific microbes, such as aerobic microbes but not anaerobes, introducing bias to the data [166]. Studies have shown that microbiome samples are stable for 2 years after being frozen at -80 °C [162,167]. Additionally, it is important to avoid multiple freeze-thaw cycles as it has been shown to affect microbial sample stability [168].

Due to the nature of the microbiome, microbiome data can be significantly affected by external factors, such as lifestyle, diet, medication and physiology. For example, for mice, housing conditions (single- or group-housed), time of cage/bedding change and fasting prior to sample collection can have significant effects on the microbial composition [169]. Additionally, technical variability is a critical issue in microbiome data analysis. Indeed, technical aspects, from DNA extraction to the choice of sequencing platform, have been found/to substantially affect data reproducibility [170,171]. Studies have also reported that the choice of DNA extraction kits, contaminants from carriers and storage methods may contribute to data variability [172–175]. Thus, standards and controls must be carefully chosen, and complete metadata should be provided along with the raw microbiome sequencing data in order to promote reproducibility and translatability of microbiome research.

5.2. Microbiome data analysis pipelines: Marker gene, metagenomics and metatranscriptomics analysis

Development of various microbiome-related experimental protocols and analytical tools have provided great opportunities in age-related microbiome research. In this section, we provide a general overview of widely used microbiome data analysis pipelines: marker gene, metagenomics and metatranscriptomics analysis (Table 2). Additionally, we discuss age-related microbiome studies that utilize each analytical method.

5.2.1. Marker gene analysis: high-level, low-resolution overview of microbial composition

Marker genes are conserved genes that contain a highly variable region, flanked by highly conserved regions that serve as primer binding sites, that can be used for detailed identification of microbial species: 16S rDNA PCR amplification is commonly used for bacteria and archaea, and ITS (internal transcribed spacer) for fungi. Marker gene analysis is well-tested, fast and cost-effective. Consequently, a significant proportion of microbiome research, not limited to age-related studies, is based on marker gene analysis data [176]. Additionally, its quantification generally correlates well with genomic content of microbial species [177–179]. However, it is important to note that marker gene data is susceptible to biases rising from variable region selection, amplicon size and number of PCR cycles [180,181]. Thus, choice of amplicon primer will have significant effect on the resolution of data [182], and it is highly recommended to review primers used in the Earth Microbiome Project [183]. Marker gene analysis has been extensively reviewed

Table 2

Pros and cons of marker gene analysis, metagenomics and metatranscriptomics.

Method	Cost	Analysis	Use in aging research	Pros	Cons
Marker gene analysis	Low	Simple	+++++	- General overview of microbial communities - Large public datasets available - High resolution (species and strain level) - Direct measure of functional microbial genes - Assemble novel microbial genomes - Measure actively transcribing microbial genes - Directly observe microbial activity changes	- Cannot discriminate among live, dead or active features - Low resolution (genus level) - Primer choice significantly affects data - Cannot discriminate among live, dead or active features - Prone to host contamination-derived errors
Metagenomics	High	Complex	++		
Meta-transcriptomics	Medium	Complex	+		- Prone to host mRNA and rRNA contamination - May fail to detect rare species or low expressed microbial genes

elsewhere [184]. PhyloChip is another 16S rRNA gene-based method for tracking microbial communities – this microarray-based technique analyzes all nine variable regions of the 16S rRNA gene [185]. In terms of aging research, PhyloChip was used in a study to analyze age-associated changes in the microbial composition in hoatzin, a South American strict folivorous bird [186].

Typically, similar sequences detected from the marker gene analyses are clustered together into Operational Taxonomic Units [OTUs]. This process, called OTU picking, consolidates similar sequences into single features and thus, merges sequence variants and may lose subtle - but real - biological sequence variants. Recent studies have started to prefer the oligotyping method to capture position-specific information from marker gene sequencing data [187]. In this method, exact sequence variants are used to distinguish between closely related, but distinct, taxa. Widely used algorithms such as Deblur and DADA2 implements this method to allow detection of subtle variations between sequences and thus enable greater sensitivity in microbiome analysis from the marker gene method [188,189]. Marker gene data can also be used to infer putative biological functions of the identified microbial community through predictive functional profiling [179,190,191]. This analysis method links the feature-abundance data from the marker gene analysis with available microbial genomes to predict the meta-genome content and biological functions. A variety of open source microbial genome references are available, including Silva, GreenGenes, and iMGMC [192–194]. It is important to note that different microbial genome references are reported to show varying degrees of sensitivity towards different microbial composition arising from specific host organism and/or sampling site [194,195]. Thus, choice of genome reference can have substantial differences on the final result (Fig. 2).

5.2.2. Metagenomics: high-resolution with genome-level information

Metagenomics is used to sequence all the microbial genomes within a given sample. This technique captures all the DNA molecules present in the sample, spanning not only bacteria, but viral and eukaryotic DNA (including that of the host). Compared to marker gene analysis, metagenomics data yields more detailed genomic information and taxonomic resolution. Additionally, this technique allows detection of microbial species to strain level and enables *de novo* metagenome assembly using short DNA sequence reads, if desired [196,197]. In addition, metagenomic studies directly enable the detection of the actual gene products present in the sample, thus giving a true window into the biological functions that the microbiome may perform [198,199]. However, it is substantially more expensive than marker gene analysis, and thus more challenging to scale up to bigger comparative studies. For a

thorough review of metagenomics analysis, refer to Refs. [200,201].

Relevant to aging, metagenomic profiling of gut microbiome from young and elderly individuals and centenarians revealed distinct characteristics of microbiome structure and function of the different age groups, identifying 116 microbial genes that significantly correlated with aging [202]. For example, the study showed that a key feature of the gut microbiome profile of centenarians is the overall increase in *Proteobacteria* and a re-arrangement in *Firmicutes* compared to young adults [202]. *Proteobacteria* has been reported to contribute to systemic inflammation [203,204]. In support of these findings, high levels of plasma interleukin-6 and interleukin-8 were detected in centenarians, although a possible pro-longevity effect of the abundant *Proteobacteria* in centenarians needs further investigation [70]. Additionally, a study by Pasolli E. et al. conducted a large-scale metagenomic analysis of human microbiome data from 9316 metagenomes spanning 46 datasets from various populations, body sites, including oral cavity, gut, skin and vagina, and host ages, spanning ages of less than 1 to over 65 [205]. Utilization of such resources will provide new insights and comprehensive understanding of the functional relationship between the microbiome and host aging.

5.2.3. Metatranscriptomics: characterization of microbial gene expression

Metatranscriptomics uses RNA-sequencing technique to profile transcription of microbiomes present in a sample. Metatranscriptomics has been argued to best represent functionality of live microbiome and thus can provide unique insights of the sample [206]. When preparing metatranscriptomics libraries, a number of considerations are required due to host RNA contamination, such as from abundant host rRNA, and preservation of RNA quality. To note, metatranscriptomics may miss the activity of rare species in the samples, due to relatively lower gene expression. For a more thorough review, refer to Ref. [207,208].

To our knowledge, there are only a limited number of metatranscriptomics studies that have been performed in the context of aging. In 2018, a large-scale investigation on 372 human male fecal metatranscriptomics was published [209]. The study involved male subjects from varying age groups, from 18 to 81 of age. Interestingly, the study revealed noticeable differences between metatranscriptomic and metagenomic data [209]. Such finding suggests the importance of multi-omics in microbiome research to accurately characterize taxonomic profiles and interpret physiological effects.

5.3. Multi-omics and multi-site analysis of the microbiome

Characterization and analysis of the microbiome have revealed immense taxonomic and genomic diversity of the microbiome and implicate more important functions of the microbiome to be




Fig. 2. General workflow for marker gene analysis.

For marker gene analysis, after sample and metadata collection, choice of primer set should be carefully considered for library preparation. Quality control of sequenced raw data, such as primer and barcode sequence removal, is strongly recommended for accurate data analysis. Features (exact sequence variants) are denoised/clustered to produce a feature-abundance matrix. Feature-abundance matrix can be used for high-level analysis, including taxonomy classification, diversity and differential abundance analysis and predictive functional profiling.

revealed in future studies. However, due to the complex functional interaction between the host and the microbiome, establishment of a causal function of the microbiome should be done with caution [210]. A multi-omics approach involving analyses of transcriptomes, proteomes, metabolomes, and immunomes along with microbiome data analysis pipelines discussed above will provide unique insights in characterizing and understanding the roles of the microbiome. Additionally, a comprehensive analysis of different microbiomes, such as skin, oral and vaginal, will be fundamental in elucidating causal functions of the microbiome on host health and longevity.

The human skin is reported to possess a distinctive microbial composition and is estimated to inhabit approximately one billion bacteria per square centimeter of skin [12,211]. Similar to other microbiomes, the skin microbiome has been shown to undergo various changes in composition throughout lifespan of the host [212]. For example, in a study of cheek microbiomes, specific genera of the *Bacteroidetes* and *Firmicutes* phyla were found on the young and specific genera of the *Actinobacteria* phyla were found only in the older age group [212]. Interestingly, Shibagaki, N. et al. Suggested that the age-associated changes in the skin microbiome is

largely influenced by the oral bacteria [213]. More specifically, microbial species that were found in greater abundance in the older age group, and thus contributed to differentiate the skin microbiomes of the different age groups, were identified as bacteria frequently found in the oral cavity, including *Streptococcus*, *Rothia* and *Veillonella* [213]. The oral microbiome has been shown to affect the whole body of the host and has been associated with a number of systemic diseases [214]. As an initiation point of digestion, the oral microbiome has been shown to impact the gut microbial composition [215]. Interestingly, the gut microbiome has been shown to affect appetite and feeding behavior of the host through the release of SCFAs [132]. Consequently, in term, the gut microbiome can affect microbial composition of the oral cavity. Together, these studies indicate a functional network among the different microbiomes on a single host and emphasize the importance of systematic investigation in microbiome research.

6. Summary and perspective

Recent availability of methodological and analytical tools has prompted researchers around the world to investigate the

functions of the microbiome and their effects on the well-being of its host. For example, a recently published study developed a human gut microbiome aging clock based on a gut metagenomics data-trained deep learning model [216]. The model was shown to achieve the mean absolute error of 5.91 years, demonstrating that generalizable indicators of age can be derived from microbiome data [216]. With increasing understanding of the importance of the gut microbiome in host longevity, it is anticipated that we will be able to identify and predict risk factors of age-onset gut dysbiosis in the near future.

In this review, we described general changes in the microbiome with age and key findings that implicate a bidirectional relationship between the host and the microbiome. We also discussed sexual dimorphism and various confounding technical factors that must be considered when analyzing and interpreting microbiome data. Over a century ago, Elie Metchnikoff hypothesized that frailty might be delayed by manipulating the gut microbiome with host-friendly bacteria found in yogurt [217]. Intriguingly, a study of yogurt consumption in Japanese individuals observed significant disparities between the two sexes in terms of induced changes to the gut microbiome, emphasizing widespread sex-dimorphism in host physiology [218]. Although it is certain that rigorous and reproducible characterization of microbiome data will offer great opportunities to develop new diagnostic biomarker in aging, key confounding (or biologically-relevant) factors, including sex-dimorphism, will need to be carefully considered. Finally, a comprehensive analysis of different microbiomes (e.g. oral, skin) and how they interact with each other, will be fundamental in elucidating causal functions of the microbiome on host health and longevity.

Funding

B.A.B is supported by NIA grants R00 AG049934 and R21 AG063739, an innovator grant from the Rose Hills foundation, and a seed grant from the NAVIGAGE foundation.

Declaration of competing interest

The authors declare that they have no conflict of interest to disclose.

Acknowledgements

We apologize to authors whose work couldn't be cited due to space constraints.

References

- [1] I.I. Metchnikoff, *The Prolongation of Life: Optimistic Studies*, Springer Publishing Company, 2004.
- [2] Integrative HMP (iHMP) Research Network Consortium, The integrative human microbiome project, *Nature* 569 (7758) (2019) 641–648.
- [3] R. Nagpal, R. Mainali, S. Ahmadi, S. Wang, R. Singh, K. Kavanagh, D.W. Kitzman, A. Kushugulova, F. Marotta, H. Yadav, Gut microbiome and aging: physiological and mechanistic insights, *Nutr. Healthy Aging* 4 (4) (2018) 267–285.
- [4] B. Bana, F. Cabreiro, The microbiome and aging, *Annu. Rev. Genet.* 53 (2019) 239–261.
- [5] F.D.D. Aleman, D.R. Valenzano, Microbiome evolution during host aging, *PLoS Pathog.* 15 (7) (2019), e1007727.
- [6] C. Xu, H. Zhu, P. Qiu, Aging progression of human gut microbiota, *BMC Microbiol.* 19 (1) (2019) 236.
- [7] J. Lederberg, A.T. McCray, Ome SweetOmics—A genealogical treasury of words, *Scientist* 15 (7) (2001), 8–8.
- [8] C. Human Microbiome Project, Structure, function and diversity of the healthy human microbiome, *Nature* 486 (7402) (2012) 207–214.
- [9] A.K. Nash, T.A. Auchtung, M.C. Wong, D.P. Smith, J.R. Gesell, M.C. Ross, C.J. Stewart, G.A. Metcalf, D.M. Muzny, R.A. Gibbs, N.J. Ajami, J.F. Petrosino, The gut mycobiome of the Human Microbiome Project healthy cohort, *Microbiome* 5 (1) (2017) 153.
- [10] J.A. Gilbert, M.J. Blaser, J.G. Caporaso, J.K. Jansson, S.V. Lynch, R. Knight, Current understanding of the human microbiome, *Nat. Med.* 24 (4) (2018) 392–400.
- [11] G.A. Cresci, E. Bawden, Gut microbiome: what we do and don't know, *Nutr. Clin. Pract.* 30 (6) (2015) 734–746.
- [12] A.L. Byrd, Y. Belkaid, J.A. Segre, The human skin microbiome, *Nat. Rev. Microbiol.* 16 (3) (2018) 143–155.
- [13] J.Y. An, K.A. Kerns, A. Ouellette, L. Robinson, H.D. Morris, C. Kaczorowski, S.I. Park, T. Mekvanich, A. Kang, J.S. McLean, T.C. Cox, M. Kaeberlein, Rapamycin rejuvenates oral health in aging mice, *elife* 9 (2020).
- [14] M.A. Curtis, P.I. Diaz, T.E. Van Dyke, The role of the microbiota in periodontal disease, *Periodontology* 83 (1) (2000) 14–25, 2020.
- [15] Buchtta, V., Vaginal microbiome. *Ceska Gynekol.* 83 (5) 371–379.
- [16] M. Li, B. Wang, M. Zhang, M. Rantalainen, S. Wang, H. Zhou, Y. Zhang, J. Shen, X. Pang, M. Zhang, H. Wei, Y. Chen, H. Lu, J. Zuo, M. Su, Y. Qiu, W. Jia, C. Xiao, L.M. Smith, S. Yang, E. Holmes, H. Tang, G. Zhao, J.K. Nicholson, L. Li, L. Zhao, Symbiotic gut microbes modulate human metabolic phenotypes, *Proc. Natl. Acad. Sci. U. S. A.* 105 (6) (2008) 2117–2122.
- [17] E.A. Eloe-Fadrosh, D.A. Rasko, The human microbiome: from symbiosis to pathogenesis, *Annu. Rev. Med.* 64 (2013) 145–163.
- [18] F. Backhed, R.E. Ley, J.L. Sonnenburg, D.A. Peterson, J.I. Gordon, Host-bacterial mutualism in the human intestine, *Science* 307 (5717) (2005) 1915–1920.
- [19] L. Dethlefsen, M. McFall-Ngai, D.A. Relman, An ecological and evolutionary perspective on human-microbe mutualism and disease, *Nature* 449 (7164) (2007) 811–818.
- [20] E. Bianconi, A. Piovesan, F. Facchini, A. Beraudi, R. Casadei, F. Frabetti, L. Vitale, M.C. Pelleri, S. Tassani, F. Piva, S. Perez-Amodio, P. Strippoli, S. Canaider, An estimation of the number of cells in the human body, *Ann. Hum. Biol.* 40 (6) (2013) 463–471.
- [21] D.C. Savage, Microbial ecology of the gastrointestinal tract, *Annu. Rev. Microbiol.* 31 (1977) 107–133.
- [22] A. Kumarappah, H. Senderovich, Therapeutic touch in the management of responsive behavior in patients with dementia, *Adv. Mind Body Med.* 30 (4) (2016) 8–13.
- [23] R. Sender, S. Fuchs, R. Milo, Are we really vastly outnumbered? Revisiting the ratio of bacterial to host cells in humans, *Cell* 164 (3) (2016) 337–340.
- [24] J.J. Dobbins, Prescott's microbiology, *J. Microbiol. Biol. Educ.: JM&BE* 11 (1) (2010) 64.
- [25] B. Lakshminarayanan, C. Stanton, P.W. O'Toole, R.P. Ross, Compositional dynamics of the human intestinal microbiota with aging: implications for health, *J. Nutr. Health Aging* 18 (9) (2014) 773–786.
- [26] N. Thevarajan, A. Puchta, C. Schulz, A. Naidoo, J.C. Szamosi, C.P. Verschoor, D. Loukov, L.P. Schenck, J. Jury, K.P. Foley, J.D. Schertzer, M.J. Larche, D.J. Davidson, E.F. Verdu, M.G. Surette, D.M.E. Bowdish, Age-associated microbial dysbiosis promotes intestinal permeability, systemic inflammation, and macrophage dysfunction, *Cell Host Microbe* 21 (4) (2017) 455–466 e454.
- [27] M.J. Claesson, I.B. Jeffery, S. Conde, S.E. Power, E.M. O'Connor, S. Cusack, H.M. Harris, M. Coakley, B. Lakshminarayanan, O. O'Sullivan, G.F. Fitzgerald, J. Deane, M. O'Connor, N. Harnedy, K. O'Connor, D. O'Mahony, D. van Sinderen, M. Wallace, L. Brennan, C. Stanton, J.R. Marchesi, A.P. Fitzgerald, F. Shanahan, C. Hill, R.P. Ross, P.W. O'Toole, Gut microbiota composition correlates with diet and health in the elderly, *Nature* 488 (7410) (2012) 178–184.
- [28] H.J. Zapata, V.J. Quagliarello, The microbiota and microbiome in aging: potential implications in health and age-related diseases, *J. Am. Geriatr. Soc.* 63 (4) (2015) 776–781.
- [29] J.K. Nicholson, E. Holmes, J. Kinross, R. Burcelin, G. Gibson, W. Jia, S. Pettersson, Host-gut microbiota metabolic interactions, *Science* 336 (6086) (2012) 1262–1267.
- [30] R. Burcelin, M. Serino, C. Chabo, V. Blasco-Baque, J. Amar, Gut microbiota and diabetes: from pathogenesis to therapeutic perspective, *Acta Diabetol.* 48 (4) (2011) 257–273.
- [31] S.M. Collins, M. Surette, P. Bercik, The interplay between the intestinal microbiota and the brain, *Nat. Rev. Microbiol.* 10 (11) (2012) 735–742.
- [32] R.A. Luna, J.A. Foster, Gut brain axis: diet microbiota interactions and implications for modulation of anxiety and depression, *Curr. Opin. Biotechnol.* 32 (2015) 35–41.
- [33] N. Salazar, L. Valdes-Varela, S. Gonzalez, M. Gueimonde, C.G. de Los Reyes-Gavilan, Nutrition and the gut microbiome in the elderly, *Gut Microb.* 8 (2) (2017) 82–97.
- [34] D. Mariat, O. Firmesse, F. Levenez, V. Guimaraes, H. Sokol, J. Dore, G. Corthier, J.P. Furet, The Firmicutes/Bacteroidetes ratio of the human microbiota changes with age, *BMC Microbiol.* 9 (2009) 123.
- [35] M.J. Hopkins, G.T. Macfarlane, Changes in predominant bacterial populations in human faeces with age and with *Clostridium difficile* infection, *J. Med. Microbiol.* 51 (5) (2002) 448–454.
- [36] E. Jimenez, L. Fernandez, M.L. Marin, R. Martin, J.M. Odriozola, C. Nueno-Palop, A. Narbad, M. Olivares, J. Xaus, J.M. Rodriguez, Isolation of commensal bacteria from umbilical cord blood of healthy neonates born by cesarean section, *Curr. Microbiol.* 51 (4) (2005) 270–274.
- [37] K. Aagaard, J. Ma, K.M. Antony, R. Ganu, J. Petrosino, J. Versalovic, The placenta harbors a unique microbiome, *Sci. Transl. Med.* 6 (237) (2014), 237ra265.

- [38] M.C. Collado, S. Rautava, J. Aakko, E. Isolauri, S. Salminen, Human gut colonisation may be initiated in utero by distinct microbial communities in the placenta and amniotic fluid, *Sci. Rep.* 6 (2016) 23129.
- [39] T. Odamaki, K. Kato, H. Sugahara, N. Hashikura, S. Takahashi, J.Z. Xiao, F. Abe, R. Osawa, Age-related changes in gut microbiota composition from newborn to centenarian: a cross-sectional study, *BMC Microbiol.* 16 (2016) 90.
- [40] N.A. Bokulich, J. Chung, T. Battaglia, N. Henderson, M. Jay, H. Li, D.L. A. F. Wu, G.I. Perez-Perez, Y. Chen, W. Schweizer, X. Zheng, M. Contreras, M.G. Dominguez-Bello, M.J. Blaser, Antibiotics, birth mode, and diet shape microbiome maturation during early life, *Sci. Transl. Med.* 8 (343) (2016), 343ra382.
- [41] F. Backhed, J. Roswall, Y. Peng, Q. Feng, H. Jia, P. Kovatcheva-Datchary, Y. Li, Y. Xia, H. Xie, H. Zhong, M.T. Khan, J. Zhang, J. Li, L. Xiao, J. Al-Aama, D. Zhang, Y.S. Lee, D. Kotowska, C. Colding, V. Tremaroli, Y. Yin, S. Bergman, X. Xu, L. Madsen, K. Kristiansen, J. Dahlgren, J. Wang, Dynamics and stabilization of the human gut microbiome during the first year of life, *Cell Host Microbe* 17 (6) (2015) 852.
- [42] M. Tanaka, J. Nakayama, Development of the gut microbiota in infancy and its impact on health in later life, *Allergol. Int.* 66 (4) (2017) 515–522.
- [43] R. Nagpal, T. Kurakawa, H. Tsuji, T. Takahashi, K. Kawashima, S. Nagata, K. Nomoto, Y. Yamashiro, Evolution of gut Bifidobacterium population in healthy Japanese infants over the first three years of life: a quantitative assessment, *Sci. Rep.* 7 (1) (2017) 10097.
- [44] R. Nagpal, H. Tsuji, T. Takahashi, K. Nomoto, K. Kawashima, S. Nagata, Y. Yamashiro, Ontogeny of the gut microbiota composition in healthy, full-term, vaginally born and breast-fed infants over the first 3 years of life: a quantitative bird's-eye view, *Front. Microbiol.* 8 (2017) 1388.
- [45] T. Yatsunenko, F.E. Rey, M.J. Manary, I. Trehan, M.G. Dominguez-Bello, M. Contreras, M. Magris, G. Hidalgo, R.N. Baldassano, A.P. Anokhin, A.C. Heath, B. Warner, J. Reeder, J. Kuczynski, J.G. Caporaso, C.A. Lozupone, C. Lauber, J.C. Clemente, D. Knights, R. Knight, J.I. Gordon, Human gut microbiome viewed across age and geography, *Nature* 486 (7402) (2012) 222–227.
- [46] C. De Filippo, D. Cavalieri, M. Di Paola, M. Ramazzotti, J.B. Poulet, S. Massart, S. Collini, G. Pieraccini, P. Lionetti, Impact of diet in shaping gut microbiota revealed by a comparative study in children from Europe and rural Africa, *Proc. Natl. Acad. Sci. U. S. A.* 107 (33) (2010) 14691–14696.
- [47] L.A. David, C.F. Maurice, R.N. Carmody, D.B. Gootenberg, J.E. Button, B.E. Wolfe, A.V. Ling, A.S. Devlin, Y. Varma, M.A. Fischbach, S.B. Biddinger, R.J. Dutton, P.J. Turnbaugh, Diet rapidly and reproducibly alters the human gut microbiome, *Nature* 505 (7484) (2014) 559–563.
- [48] E.B. Hollister, C. Gao, J. Versalovic, Compositional and functional features of the gastrointestinal microbiome and their effects on human health, *Gastroenterology* 146 (6) (2014) 1449–1458.
- [49] P.B. Eckburg, E.M. Bik, C.N. Bernstein, E. Purdom, L. Dethlefsen, M. Sargent, S.R. Gill, K.E. Nelson, D.A. Relman, Diversity of the human intestinal microbial flora, *Science* 308 (5728) (2005) 1635–1638.
- [50] E.M. Quigley, Gut bacteria in health and disease, *Gastroenterol. Hepatol.* 9 (9) (2013) 560–569.
- [51] H.M. Wexler, Bacteroides: the good, the bad, and the nitty-gritty, *Clin. Microbiol. Rev.* 20 (4) (2007) 593–621.
- [52] G. den Besten, K. van Eunen, A.K. Groen, K. Venema, D.-J. Reijngoud, B.M. Bakker, The role of short-chain fatty acids in the interplay between diet, gut microbiota, and host energy metabolism, *J. Lipid Res.* 54 (9) (2013) 2325–2340.
- [53] P. Louis, S.H. Duncan, S.I. McCrae, J. Millar, M.S. Jackson, H.J. Flint, Restricted distribution of the butyrate kinase pathway among butyrate-producing bacteria from the human colon, *J. Bacteriol.* 186 (7) (2004) 2099–2106.
- [54] P. Louis, P. Young, G. Holtrop, H.J. Flint, Diversity of human colonic butyrate-producing bacteria revealed by analysis of the butyryl-CoA:acetate CoA-transferase gene, *Environ. Microbiol.* 12 (2) (2010) 304–314.
- [55] L.V. Hooper, T. Midtvedt, J.I. Gordon, How host-microbial interactions shape the nutrient environment of the mammalian intestine, *Annu. Rev. Nutr.* 22 (2002) 283–307.
- [56] A.C. Razavi, K.S. Potts, T.N. Kelly, L.A. Bazzano, Sex, gut microbiome, and cardiovascular disease risk, *Biol. Sex Differ.* 10 (1) (2019) 29.
- [57] C. Indiani, K.F. Rizzardi, P.M. Castelo, L.F.C. Ferraz, M. Darrieux, T.M. Parisotto, Childhood obesity and firmicutes/bacteroidetes ratio in the gut microbiota: a systematic review, *Child. Obes.* 14 (8) (2018) 501–509.
- [58] R.E. Ley, P.J. Turnbaugh, S. Klein, J.I. Gordon, Microbial ecology: human gut microbes associated with obesity, *Nature* 444 (7122) (2006) 1022–1023.
- [59] E. Rinninella, P. Raoul, M. Cintoni, F. Franceschi, G.A.D. Miggiano, A. Gasbarrini, M.C. Mele, What is the healthy gut microbiota composition? A changing ecosystem across age, environment, diet, and diseases, *Microorganisms* 7 (1) (2019).
- [60] A. Koliada, G. Syzenko, V. Moseiko, L. Budovska, K. Puchkov, V. Perederiy, Y. Gavalko, A. Dorofeyev, M. Romanenko, S. Tkach, L. Sineok, O. Lushchak, A. Vaiserman, Association between body mass index and Firmicutes/Bacteroidetes ratio in an adult Ukrainian population, *BMC Microbiol.* 17 (1) (2017) 120.
- [61] M.J. Claesson, S. Cusack, O. O'Sullivan, R. Greene-Diniz, H. de Weerd, E. Flannery, J.R. Marchesi, D. Falush, T. Dinan, G. Fitzgerald, C. Stanton, D. van Sinderen, M. O'Connor, N. Harnedy, K. O'Connor, C. Henry, D. O'Mahony, A.P. Fitzgerald, F. Shanahan, C. Twomey, C. Hill, R.P. Ross, P.W. O'Toole, Composition, variability, and temporal stability of the intestinal microbiota of the elderly, *Proc. Natl. Acad. Sci. U. S. A.* 108 (Suppl 1) (2011) 4586–4591.
- [62] A.D. Kostic, D. Gevers, H. Siljander, T. Vatanen, T. Hyotylainen, A.M. Hamalainen, A. Peet, V. Tillmann, P. Poho, I. Mattila, H. Lahdesmaki, E.A. Franzosa, O. Vaarala, M. de Goffau, H. Harmsen, J. Ilonen, S.M. Virtanen, C.B. Clish, M. Oresic, C. Huttenhower, M. Knip, D.S. Group, R.J. Xavier, The dynamics of the human infant gut microbiome in development and in progression toward type 1 diabetes, *Cell Host Microbe* 17 (2) (2015) 260–273.
- [63] J.U. Scher, A. Sczesnak, R.S. Longman, N. Segata, C. Ubeda, C. Bielski, T. Rostron, V. Cerundolo, E.G. Palmer, S.B. Abramson, C. Huttenhower, D.R. Littman, Expansion of intestinal Prevotella copri correlates with enhanced susceptibility to arthritis, *eLife* 2 (2013), e01202.
- [64] A.R. Moschen, R.R. Gerner, J. Wang, V. Klepsch, T.E. Adolph, S.J. Reider, H. Hackl, A. Pfister, J. Schilling, P.L. Moser, S.L. Kempster, A. Swidsinski, D. Orth Holler, G. Weiss, J.F. Baines, A. Kaser, H. Tilg, Lipocalin 2 protects from inflammation and tumorigenesis associated with gut microbiota alterations, *Cell Host Microbe* 19 (4) (2016) 455–469.
- [65] P.W. O'Toole, I.B. Jeffery, Gut microbiota and aging, *Science* 350 (6265) (2015) 1214–1215.
- [66] F. Gavini, C. Cayuela, J.-M. Antoine, C. Lecocq, B. Lefebvre, J.-M. Membré, C. Neut, Differences in the distribution of bifidobacterial and enterobacterial species in human faecal microflora of three different (children, adults, elderly) age groups, *Microb. Ecol. Health Dis.* 13 (1) (2001) 40–45.
- [67] S. Mueller, K. Saunier, C. Hanisch, E. Norin, L. Alm, T. Midtvedt, A. Cresci, S. Silvi, C. Orpianesi, M.C. Verdenelli, T. Clavel, C. Koebnick, H.J. Zunft, J. Dore, M. Blaut, Differences in fecal microbiota in different European study populations in relation to age, gender, and country: a cross-sectional study, *Appl. Environ. Microbiol.* 72 (2) (2006) 1027–1033.
- [68] Y. Benno, K. Endo, T. Mizutani, Y. Namba, T. Komori, T. Mitsuoka, Comparison of fecal microflora of elderly persons in rural and urban areas of Japan, *Appl. Environ. Microbiol.* 55 (5) (1989) 1100–1105.
- [69] E. Biagi, C. Franceschi, S. Rampelli, M. Severgnini, R. Ostan, S. Turroni, C. Consolandi, S. Quercia, M. Scurti, D. Monti, M. Capri, P. Brigidi, M. Candela, Gut microbiota and extreme longevity, *Curr. Biol.* 26 (11) (2016) 1480–1485.
- [70] E. Biagi, L. Nylund, M. Candela, R. Ostan, L. Bucci, E. Pini, J. Nikkila, D. Monti, R. Satokari, C. Franceschi, P. Brigidi, W. De Vos, Through ageing, and beyond: gut microbiota and inflammatory status in seniors and centenarians, *PLoS One* 5 (5) (2010), e10667.
- [71] T. Komura, C. Yasui, H. Miyamoto, Y. Nishikawa, Caenorhabditis elegans as an alternative model host for legionella pneumophila, and protective effects of *Bifidobacterium infantis*, *Appl. Environ. Microbiol.* 76 (12) (2010) 4105–4108.
- [72] T. Komura, T. Ikeda, C. Yasui, S. Saeki, Y. Nishikawa, Mechanism underlying prolongevity induced by bifidobacteria in *Caenorhabditis elegans*, *Bio-gerontology* 14 (1) (2013) 73–87.
- [73] J.K. Goodrich, J.L. Waters, A.C. Poole, J.L. Sutter, O. Koren, R. Blekhman, M. Beaumont, W. Van Treuren, R. Knight, J.T. Bell, T.D. Spector, A.G. Clark, R.E. Ley, Human genetics shape the gut microbiome, *Cell* 159 (4) (2014) 789–799.
- [74] R.I. Clark, A. Salazar, R. Yamada, S. Fitz-Gibbon, M. Morselli, J. Alcaraz, A. Rana, M. Rera, M. Pellegrini, W.W. Ja, D.W. Walker, Distinct shifts in microbiota composition during *Drosophila* aging impair intestinal function and drive mortality, *Cell Rep.* 12 (10) (2015) 1656–1667.
- [75] M. Rera, R.I. Clark, D.W. Walker, Intestinal barrier dysfunction links metabolic and inflammatory markers of aging to death in *Drosophila*, *Proc. Natl. Acad. Sci. U. S. A.* 109 (52) (2012) 21528–21533.
- [76] P. Smith, D. Willemse, M. Popkes, F. Metge, E. Gandiwa, M. Reichard, D.R. Valenzano, Regulation of life span by the gut microbiota in the short-lived African turquoise killifish, *eLife* 6 (2017).
- [77] F. Cabreiro, D. Gems, Worms need microbes too: microbiota, health and aging in *Caenorhabditis elegans*, *EMBO Mol. Med.* 5 (9) (2013) 1300–1310.
- [78] S.P. van Tongeren, J.P. Slaets, H.J. Harmsen, G.W. Welling, Fecal microbiota composition and frailty, *Appl. Environ. Microbiol.* 71 (10) (2005) 6438–6442.
- [79] F.A. Carvalho, O. Koren, J.K. Goodrich, M.E. Johansson, I. Nalbantoglu, J.D. Aitken, Y. Su, B. Chassaing, W.A. Walters, A. Gonzalez, J.C. Clemente, T.C. Cullender, N. Barnich, A. Darfeuille-Michaud, M. Vijay-Kumar, R. Knight, R.E. Ley, A.T. Gewirtz, Transient inability to manage proteobacteria promotes chronic gut inflammation in TLR5-deficient mice, *Cell Host Microbe* 12 (2) (2012) 139–152.
- [80] M.G. Langille, C.J. Meehan, J.E. Koenig, A.S. Dhanani, R.A. Rose, S.E. Howlett, R.G. Beiko, Microbial shifts in the aging mouse gut, *Microbiome* 2 (1) (2014) 50.
- [81] B. Flemer, N. Gaci, G. Borrel, I.R. Sanderson, P.P. Chaudhary, W. Tottey, P.W. O'Toole, J.F. Brugere, Fecal microbiota variation across the lifespan of the healthy laboratory rat, *Gut Microb.* 8 (5) (2017) 428–439.
- [82] C. Barcena, R. Valdes-Mas, P. Mayoral, C. Garabaya, S. Durand, F. Rodriguez, M.T. Fernandez-Garcia, N. Salazar, A.M. Nogacka, N. Garatachea, N. Bossut, F. Aprahamian, A. Lucia, G. Kroemer, J.M.P. Freije, P.M. Quiros, C. Lopez-Otin, Healthspan and lifespan extension by fecal microbiota transplantation into progeroid mice, *Nat. Med.* 25 (8) (2019) 1234–1242.
- [83] S. Wang, M. Huang, X. You, J. Zhao, L. Chen, L. Wang, Y. Luo, Y. Chen, Gut microbiota mediates the anti-obesity effect of calorie restriction in mice, *Sci. Rep.* 8 (1) (2018) 13037.
- [84] F. Pan, L. Zhang, M. Li, Y. Hu, B. Zeng, H. Yuan, L. Zhao, C. Zhang, Predominant gut *Lactobacillus murinus* strain mediates anti-inflammatory effects in calorie-restricted mice, *Microbiome* 6 (1) (2018) 54.

- [85] R. Blumberg, F. Powrie, Microbiota, disease, and back to health: a metastable journey, *Sci. Transl. Med.* 4 (137) (2012) 137rv137.
- [86] S. Carding, K. Verbeke, D.T. Vipond, B.M. Corfe, L.J. Owen, Dysbiosis of the gut microbiota in disease, *Microb. Ecol. Health Dis.* 26 (2015) 26191.
- [87] M.T. Khan, M. Nieuwdorp, F. Backhed, Microbial modulation of insulin sensitivity, *Cell Metabol.* 20 (5) (2014) 753–760.
- [88] C. Franceschi, J. Campisi, Chronic inflammation (inflammaging) and its potential contribution to age-associated diseases, *J. Gerontol. A Biol. Sci. Med. Sci.* 69 (Suppl 1) (2014) S4–S9.
- [89] C. Franceschi, M. Bonafe, S. Valensin, F. Olivieri, M. De Luca, E. Ottaviani, G. De Benedictis, Inflamm-aging. An evolutionary perspective on immunosenescence, *Ann. N. Y. Acad. Sci.* 908 (2000) 244–254.
- [90] A.J. Macpherson, N.L. Harris, Interactions between commensal intestinal bacteria and the immune system, *Nat. Rev. Immunol.* 4 (6) (2004) 478–485.
- [91] P. Van den Abbeele, T. Van de Wiele, W. Verstraete, S. Possemiers, The host selects mucosal and luminal associations of coevolved gut microorganisms: a novel concept, *FEMS Microbiol. Rev.* 35 (4) (2011) 681–704.
- [92] J.R. Turner, Intestinal mucosal barrier function in health and disease, *Nat. Rev. Immunol.* 9 (11) (2009) 799–809.
- [93] J. Pott, M. Hornef, Innate immune signalling at the intestinal epithelium in homeostasis and disease, *EMBO Rep.* 13 (8) (2012) 684–698.
- [94] Y. Belkaid, S. Naik, Compartmentalized and systemic control of tissue immunity by commensals, *Nat. Immunol.* 14 (7) (2013) 646–653.
- [95] M.C. Abt, L.C. Osborne, L.A. Monticelli, T.A. Doering, T. Alenghat, G.F. Sonnenberg, M.A. Paley, M. Antenus, K.L. Williams, J. Erikson, E.J. Wherry, D. Artis, Commensal bacteria calibrate the activation threshold of innate antiviral immunity, *Immunity* 37 (1) (2012) 158–170.
- [96] B.B. Yoo, S.K. Mazmanian, The enteric network: interactions between the immune and nervous systems of the gut, *Immunity* 46 (6) (2017) 910–926.
- [97] S.M. Man, Q. Zhu, L. Zhu, Z. Liu, R. Karki, A. Malik, D. Sharma, L. Li, R.K. Malireddi, P. Gurung, G. Neale, S.R. Olsen, R.A. Carter, D.J. McGoldrick, G. Wu, D. Finkelstein, P. Vogel, R.J. Gilbertson, T.D. Kanneganti, Critical role for the DNA sensor AIM2 in stem cell proliferation and cancer, *Cell* 162 (1) (2015) 45–58.
- [98] L. Guo, J. Karpac, S.L. Tran, H. Jasper, PGRP-SC2 promotes gut immune homeostasis to limit commensal dysbiosis and extend lifespan, *Cell* 156 (1–2) (2014) 109–122.
- [99] M. Levy, A.A. Kolodziejczyk, C.A. Thaiss, E. Elinav, Dysbiosis and the immune system, *Nat. Rev. Immunol.* 17 (4) (2017) 219–232.
- [100] S. Bartosch, A. Fite, G.T. Macfarlane, M.E. McMurdo, Characterization of bacterial communities in feces from healthy elderly volunteers and hospitalized elderly patients by using real-time PCR and effects of antibiotic treatment on the fecal microbiota, *Appl. Environ. Microbiol.* 70 (6) (2004) 3575–3581.
- [101] M.A. Jackson, I.B. Jeffery, M. Beaumont, J.T. Bell, A.G. Clark, R.E. Ley, P.W. O'Toole, T.D. Spector, C.J. Steves, Signatures of early frailty in the gut microbiota, *Genome Med.* 8 (1) (2016) 8.
- [102] A. Kumar, H. Wu, L.S. Collier-Hyams, J.M. Hansen, T. Li, K. Yamoah, Z.Q. Pan, D.P. Jones, A.S. Neish, Commensal bacteria modulate cullin-dependent signaling via generation of reactive oxygen species, *EMBO J.* 26 (21) (2007) 4457–4466.
- [103] E.G. Pamer, Immune responses to commensal and environmental microbes, *Nat. Immunol.* 8 (11) (2007) 1173–1178.
- [104] J. Karczewski, B. Poniedzialek, Z. Adamski, P. Rzymski, The effects of the microbiota on the host immune system, *Autoimmunity* 47 (8) (2014) 494–504.
- [105] Y. Belkaid, T.W. Hand, Role of the microbiota in immunity and inflammation, *Cell* 157 (1) (2014) 121–141.
- [106] S. Rakoff-Nahoum, J. Paglino, F. Eslami-Varzaneh, S. Edberg, R. Medzhitov, Recognition of commensal microflora by toll-like receptors is required for intestinal homeostasis, *Cell* 118 (2) (2004) 229–241.
- [107] A. Khosravi, A. Yaneez, J.G. Price, A. Chow, M. Merad, H.S. Goodridge, S.K. Mazmanian, Gut microbiota promote hematopoiesis to control bacterial infection, *Cell Host Microbe* 15 (3) (2014) 374–381.
- [108] S.K. Mazmanian, C.H. Liu, A.O. Tzianabos, D.L. Kasper, An immunomodulatory molecule of symbiotic bacteria directs maturation of the host immune system, *Cell* 122 (1) (2005) 107–118.
- [109] J.M. Mullin, M.C. Valenzano, J.J. Verrecchio, R. Kothari, Age- and diet-related increase in transepithelial colon permeability of Fischer 344 rats, *Dig. Dis. Sci.* 47 (10) (2002) 2262–2270.
- [110] P. Annaert, J. Brouwers, A. Bijnens, F. Lammert, J. Tack, P. Augustijns, Ex vivo permeability experiments in excised rat intestinal tissue and in vitro solubility measurements in aspirated human intestinal fluids support age-dependent oral drug absorption, *Eur. J. Pharmaceut. Sci.* 39 (1–3) (2010) 15–22.
- [111] S. Gelino, J.T. Chang, C. Kumsta, X. She, A. Davis, C. Nguyen, S. Panowski, M. Hansen, Intestinal autophagy improves healthspan and longevity in *C. elegans* during dietary restriction, *PLoS Genet.* 12 (7) (2016), e1006135.
- [112] J. Amar, C. Chabo, A. Waget, P. Klopp, C. Vachoux, L.G. Bermudez-Humaran, N. Smirnova, M. Berge, T. Sulpice, S. Lahtinen, A. Ouwehand, P. Langella, N. Rautonen, P.J. Sansonetti, R. Burcelin, Intestinal mucosal adherence and translocation of commensal bacteria at the early onset of type 2 diabetes: molecular mechanisms and probiotic treatment, *EMBO Mol. Med.* 3 (9) (2011) 559–572.
- [113] H.E. Raybould, Gut microbiota, epithelial function and derangements in obesity, *J. Physiol.* 590 (3) (2012) 441–446.
- [114] J. Sato, A. Kanazawa, F. Ikeda, T. Yoshihara, H. Goto, H. Abe, K. Komiya, M. Kawaguchi, T. Shimizu, T. Ogihara, Y. Tamura, Y. Sakurai, R. Yamamoto, T. Mita, Y. Fujitani, H. Fukuda, K. Nomoto, T. Takahashi, T. Asahara, T. Hirose, S. Nagata, Y. Yamashiro, H. Watada, Gut dysbiosis and detection of "live gut bacteria" in blood of Japanese patients with type 2 diabetes, *Diabetes Care* 37 (8) (2014) 2343–2350.
- [115] R.M. Al-Sadi, T.Y. Ma, IL-1 β causes an increase in intestinal epithelial tight junction permeability, *J. Immunol.* 178 (7) (2007) 4641–4649.
- [116] A.L. Man, N. Gicheva, C. Nicoletti, The impact of ageing on the intestinal epithelial barrier and immune system, *Cell. Immunol.* 289 (1–2) (2014) 112–118.
- [117] C.J. Kelly, L. Zheng, E.L. Campbell, B. Saeedi, C.C. Scholz, A.J. Bayless, K.E. Wilson, L.E. Glover, D.J. Kominsky, A. Magnuson, T.L. Weir, S.F. Ehrentraut, C. Pickel, K.A. Kuhn, J.M. Lanis, V. Nguyen, C.T. Taylor, S.P. Colgan, Crosstalk between microbiota-derived short-chain fatty acids and intestinal epithelial HIF augments tissue barrier function, *Cell Host Microbe* 17 (5) (2015) 662–671.
- [118] K. Tiilohon, A.C. Ouwehand, N. Rautonen, Human intestinal microbiota and healthy ageing, *Ageing Res. Rev.* 9 (2) (2010) 107–116.
- [119] T. Hoverstad, T. Midtvedt, Short-chain fatty acids in germfree mice and rats, *J. Nutr.* 116 (9) (1986) 1772–1776.
- [120] J.L. Fachi, J.S. Felipe, L.P. Pral, B.K. da Silva, R.O. Correa, M.C.P. de Andrade, D.M. da Fonseca, P.J. Basso, N.O.S. Camara, E.S.E.L. de Sales, F. Dos Santos Martins, S.E.S. Guima, A.M. Thomas, J.C. Setubal, Y.T. Magalhaes, F.L. Forti, T. Candreva, H.G. Rodrigues, M.B. de Jesus, S.R. Consonni, A.D.S. Farias, P. Varga-Weisz, M.A.R. Vinolo, Butyrate protects mice from Clostridium difficile-induced colitis through an HIF-1-Dependent mechanism, *Cell Rep.* 27 (3) (2019) 750–761 e757.
- [121] D. Erny, A.L. Hrabé de Angelis, D. Jaitin, P. Wieghofer, O. Staszewski, E. David, H. Keren-Shaul, T. Mahlakovic, K. Jakobshagen, T. Buch, V. Schwierzeck, O. Utermöhlen, E. Chun, W.S. Garrett, K.D. McCoy, A. Dieffenbach, P. Staeheli, B. Stecher, I. Amit, M. Prinz, Host microbiota constantly control maturation and function of microglia in the CNS, *Nat. Neurosci.* 18 (7) (2015) 965–977.
- [122] A.D. Vincent, X.Y. Wang, S.P. Parsons, W.I. Khan, J.D. Huizinga, Abnormal absorptive colonic motor activity in germ-free mice is rectified by butyrate, an effect possibly mediated by mucosal serotonin, *Am. J. Physiol. Gastrointest. Liver Physiol.* 315 (5) (2018) G896–G907.
- [123] V. Braniste, M. Al-Asmakh, C. Kowal, F. Anuar, A. Abbaspour, M. Toth, A. Korecka, N. Bakoczevic, L.G. Ng, P. Kundu, B. Gulyas, C. Halldin, K. Hultenby, H. Nilsson, H. Hebert, B.T. Volpe, B. Diamond, S. Pettersson, The gut microbiota influences blood-brain barrier permeability in mice, *Sci. Transl. Med.* 6 (263) (2014), 263ra158.
- [124] G.E. Kaiko, S.H. Ryu, O.I. Koues, P.L. Collins, L. Solnica-Krezel, E.J. Pearce, E.L. Pearce, E.M. Oltz, T.S. Stappenbeck, The colonic crypt protects stem cells from microbiota-derived metabolites, *Cell* 167 (4) (2016) 1137.
- [125] B.C.E. Peck, M.T. Shanahan, A.P. Singh, P. Sethupathy, Gut microbial influences on the mammalian intestinal stem cell niche, *Stem Cell. Int.* (2017) 2017, 5604727.
- [126] A. Koh, F. Backhed, From association to causality: the role of the gut microbiota and its functional products on host metabolism, *Mol. Cell.* 78 (4) (2020) 584–596.
- [127] Z. Shan, T. Sun, H. Huang, S. Chen, L. Chen, C. Luo, W. Yang, X. Yang, P. Yao, J. Cheng, Association between microbiota-dependent metabolite trimethylamine-N-oxide and type 2 diabetes, *Am. J. Clin. Nutr.* 106 (3) (2017) 888–894.
- [128] W.H. Tang, T. Kitai, S.L. Hazen, Gut microbiota in cardiovascular health and disease, *Circ. Res.* 120 (7) (2017) 1183–1196.
- [129] P.V. Chang, L. Hao, S. Offermanns, R. Medzhitov, The microbial metabolite butyrate regulates intestinal macrophage function via histone deacetylase inhibition, *Proc. Natl. Acad. Sci. U. S. A.* 111 (6) (2014) 2247–2252.
- [130] Y. Qin, P.A. Wade, Crosstalk between the microbiome and epigenome: messages from bugs, *J. Biochem.* 163 (2) (2018) 105–112.
- [131] K.G. Jameson, C.A. Olson, S.A. Kazmi, E.Y. Hsiao, Toward understanding microbiome-neuronal signaling, *Mol. Cell.* 78 (4) (2020) 577–583.
- [132] C.S. Byrne, E.S. Chambers, D.J. Morrison, G. Frost, The role of short chain fatty acids in appetite regulation and energy homeostasis, *Int. J. Obes.* 39 (9) (2015) 1331–1338.
- [133] N.K. Sampathkumar, J.I. Bravo, Y. Chen, P.S. Danthi, E.K. Donahue, R.W. Lai, R. Lu, L.T. Randall, N. Vinson, B.A. Benayoun, Widespread sex dimorphism in aging and age-related diseases, *Hum. Genet.* 139 (3) (2020) 333–356.
- [134] R. Ostan, D. Monti, P. Gueresi, M. Bussolotto, C. Franceschi, G. Baggio, Gender, aging and longevity in humans: an update of an intriguing/neglected scenario paving the way to a gender-specific medicine, *Clin. Sci. (Lond.)* 130 (19) (2016) 1711–1725.
- [135] K.N. Manolopoulos, F. Karpe, K.N. Frayn, Gluteofemoral body fat as a determinant of metabolic health, *Int. J. Obes.* 34 (6) (2010) 949–959.
- [136] E. Rose, M. Gedela, N. Miller, P.L. Carpenter, Pregnancy-related spontaneous coronary artery dissection: a case series and literature review, *J. Emerg. Med.* 52 (6) (2017) 867–874.
- [137] J.G. Markle, D.N. Frank, S. Morton-Toth, C.E. Robertson, L.M. Feazel, U. Rolle-Kampczyk, M. von Bergen, K.D. McCoy, A.J. Macpherson, J.S. Danska, Sex differences in the gut microbiome drive hormone-dependent regulation of autoimmunity, *Science* 339 (6123) (2013) 1084–1088.
- [138] W.T. Steegenga, M. Mischke, C. Lute, M.V. Boekschen, M.G. Pruis,

- A. Lendvai, H.J. Verkade, J. Boekhorst, H.M. Timmerman, T. Plosch, M. Muller, Sexually dimorphic characteristics of the small intestine and colon of pre-pubescent C57BL/6 mice, *Biol. Sex Differ.* 5 (2014) 11.
- [139] C. Dominianni, R. Sinha, J.J. Goedert, Z. Pei, L. Yang, R.B. Hayes, J. Ahn, Sex, body mass index, and dietary fiber intake influence the human gut microbiome, *PLoS One* 10 (4) (2015), e0124599.
- [140] C. Haro, O.A. Rangel-Zuniga, J.F. Alcalá-Díaz, F. Gómez-Delgado, P. Pérez-Martínez, J. Delgado-Lista, G.M. Quintana-Navarro, B.B. Landa, J.A. Navas-Cortes, M. Tena-Sempere, J.C. Clemente, J. López-Miranda, F. Pérez-Jiménez, A. Camargo, Intestinal microbiota is influenced by gender and body mass index, *PLoS One* 11 (5) (2016), e0154090.
- [141] L. Yurkovetskiy, M. Burrows, A.A. Khan, L. Graham, P. Volchkov, L. Becker, D. Antonopoulos, Y. Umesaki, A.V. Chervonsky, Gender bias in autoimmunity is influenced by microbiota, *Immunity* 39 (2) (2013) 400–412.
- [142] B.D. Weger, C. Gobet, J. Yeung, E. Martin, S. Jimenez, B. Betrisey, F. Foata, B. Berger, A. Balvay, A. Foussier, A. Charpagne, B. Boizet-Bonhoure, C.J. Chou, F. Naef, F. Gachon, The mouse microbiome is required for sex-specific diurnal rhythms of gene expression and metabolism, *Cell Metabol.* 29 (2) (2019) 362–382 e368.
- [143] A. Baars, A. Oosting, M. Lohuis, M. Koehorst, S. El Aidy, F. Hugenholtz, H. Smidt, M. Mischke, M.V. Boekschoten, H.J. Verkade, J. Garssen, E.M. van der Beek, J. Knol, P. de Vos, J. van Bergenhenegouwen, F. Fransen, Sex differences in lipid metabolism are affected by presence of the gut microbiota, *Sci. Rep.* 8 (1) (2018) 13426.
- [144] M.S. Thion, D. Low, A. Silvin, J. Chen, P. Grisel, J. Schulte-Schrepping, R. Blecher, T. Ulas, P. Squarzoni, G. Hoeffel, F. Couplier, E. Siopi, F.S. David, C. Scholz, F. Shihui, J. Lum, A.A. Amoyo, A. Larbi, M. Poidinger, A. Buttgererit, P.M. Lledo, M. Greter, J.K.Y. Chan, I. Amit, M. Beyer, J.L. Schultz, A. Schlitzer, S. Pettersson, F. Ginhoux, S. Garel, Microbiome influences prenatal and adult microglia in a sex-specific manner, *Cell* 172 (3) (2018) 500–516 e516.
- [145] H.B. Dodiya, T. Kuntz, S.M. Shaik, C. Baufeld, J. Leibowitz, X. Zhang, N. Gottel, X. Zhang, O. Butovsky, J.A. Gilbert, S.S. Sisodia, Sex-specific effects of microbiome perturbations on cerebral Abeta amyloidosis and microglia phenotypes, *J. Exp. Med.* 216 (7) (2019) 1542–1560.
- [146] H. Adlercreutz, F. Martin, P. Jarvenpaa, T. Fotsis, Steroid absorption and enterohepatic recycling, *Contraception* 20 (3) (1979) 201–223.
- [147] H. Eriksson, J.A. Gustafsson, Steroids in germfree and conventional rats. Distribution and excretion of labelled pregnenolone and corticosterone in male and female rats, *Eur. J. Biochem.* 15 (1) (1970) 132–139.
- [148] P. Lombardi, B. Goldin, E. Boutin, S.I. Gorbach, Metabolism of androgens and estrogens by human fecal microorganisms, *J. Steroid Biochem.* 9 (8) (1978) 795–801.
- [149] C.S. Plotell, M.J. Blaser, Microbiome and malignancy, *Cell Host Microbe* 10 (4) (2011) 324–335.
- [150] J.M. Baker, L. Al-Nakkash, M.M. Herbst-Kralovetz, Estrogen-gut microbiome axis: physiological and clinical implications, *Maturitas* 103 (2017) 45–53.
- [151] S.M. Ervin, H. Li, L. Lim, L.R. Roberts, X. Liang, S. Mani, M.R. Redinbo, Gut microbial beta-glucuronidases reactivate estrogens as components of the estrobolome that reactivate estrogens, *J. Biol. Chem.* 294 (49) (2019) 18586–18599.
- [152] R. Vemuri, K.E. Sylvia, S.L. Klein, S.C. Forster, M. Plebanski, R. Eri, K.L. Flanagan, The microgenderome revealed: sex differences in bidirectional interactions between the microbiota, hormones, immunity and disease susceptibility, *Semin. Immunopathol.* 41 (2) (2019) 265–275.
- [153] K. Kaliannan, R.C. Robertson, K. Murphy, C. Stanton, C. Kang, B. Wang, L. Hao, A.K. Bhan, J.X. Kang, Estrogen-mediated gut microbiome alterations influence sexual dimorphism in metabolic syndrome in mice, *Microbiome* 6 (1) (2018) 205.
- [154] M.H. Fraher, P.W. O'Toole, E.M. Quigley, Techniques used to characterize the gut microbiota: a guide for the clinician, *Nat. Rev. Gastroenterol. Hepatol.* 9 (6) (2012) 312–322.
- [155] P.J. Turnbaugh, R.E. Ley, M.A. Mahowald, V. Magrini, E.R. Mardis, J.I. Gordon, An obesity-associated gut microbiome with increased capacity for energy harvest, *Nature* 444 (7122) (2006) 1027–1031.
- [156] M.I. Smith, T. Yatsunenko, M.J. Manary, I. Trehan, R. Mkakosya, J. Cheng, A.L. Kau, S.S. Rich, P. Concannon, J.C. Mychaleckyj, J. Liu, E. Houpt, J.V. Li, E. Holmes, J. Nicholson, D. Knights, L.K. Ursell, R. Knight, J.I. Gordon, Gut microbiomes of Malawian twin pairs discordant for kwashiorkor, *Science* 339 (6119) (2013) 548–554.
- [157] B. Gustafsson, Germ-free rearing of rats, *Acta Anat.* 2 (3–4) (1946) 376–391.
- [158] T. Singer-Englar, G. Barlow, R. Mathur, Obesity, diabetes, and the gut microbiome: an updated review, *Expert Rev. Gastroenterol. Hepatol.* 13 (1) (2019) 3–15.
- [159] W.H. Tang, S.L. Hazen, The gut microbiome and its role in cardiovascular diseases, *Circulation* 135 (11) (2017) 1008–1010.
- [160] E. Elinav, W.S. Garrett, G. Trinchieri, J. Wargo, The cancer microbiome, *Nat. Rev. Canc.* 19 (7) (2019) 371–376.
- [161] T. Harach, N. Marungruang, N. Duthilleul, V. Cheatham, K.D. Mc Coy, G. Frisoni, J.J. Neher, F. Fak, M. Jucker, T. Lasser, T. Bolmont, Reduction of Abeta amyloid pathology in APPPS1 transgenic mice in the absence of gut microbiota, *Sci. Rep.* 7 (2017) 41802.
- [162] R. Kumar, P. Eipers, R.B. Little, M. Crowley, D.K. Crossman, E.J. Lefkowitz, C.D. Morrow, Getting started with microbiome analysis: sample acquisition to bioinformatics, *Curr. Protoc. Hum. Genet.* 82 18 (2014) 18 11–29.
- [163] C. Human Microbiome Project, A framework for human microbiome research, *Nature* 486 (7402) (2012) 215–221.
- [164] E. Vogtmann, J. Chen, M.G. Kibriya, Y. Chen, T. Islam, M. Eunes, A. Ahmed, J. Naher, A. Rahman, A. Amir, J. Shi, C.C. Abnet, H. Nelson, R. Knight, N. Chia, H. Ahsan, R. Sinha, Comparison of fecal collection methods for microbiota studies in Bangladesh, *Appl. Environ. Microbiol.* 83 (10) (2017).
- [165] M. Wasfy, B. Oyofo, A. Elgindy, A. Churilla, Comparison of preservation media for storage of stool samples, *J. Clin. Microbiol.* 33 (8) (1995) 2176–2178.
- [166] J.M. Nechvatal, J.L. Ram, M.D. Basson, P. Namprachan, S.R. Niec, K.Z. Badsha, L.H. Matherly, A.P. Majumdar, I. Kato, Fecal collection, ambient preservation, and DNA extraction for PCR amplification of bacterial and human markers from human feces, *J. Microbiol. Methods* 72 (2) (2008) 124–132.
- [167] A.G. Shaw, K. Sim, E. Powell, E. Cornwell, T. Cramer, Z.E. McClure, M.S. Li, J.S. Kroll, Latitude in sample handling and storage for infant faecal microbiota studies: the elephant in the room? *Microbiome* 4 (1) (2016) 40.
- [168] V. Thomas, J. Clark, J. Dore, Fecal microbiota analysis: an overview of sample collection methods and sequencing strategies, *Future Microbiol.* 10 (9) (2015) 1485–1504.
- [169] A.C. Ericsson, J. Gagliardi, D. Bouhan, W.G. Spollen, S.A. Givan, C.L. Franklin, The influence of caging, bedding, and diet on the composition of the microbiota in different regions of the mouse gut, *Sci. Rep.* 8 (1) (2018) 4065.
- [170] R. Sinha, G. Abu-Ali, E. Vogtmann, A.A. Fodor, B. Ren, A. Amir, E. Schwager, J. Crabtree, S. Ma, C. Microbiome Quality Control Project, C.C. Abnet, R. Knight, O. White, C. Huttenhower, Assessment of variation in microbial community amplicon sequencing by the Microbiome Quality Control (MBQC) project consortium, *Nat. Biotechnol.* 35 (11) (2017) 1077–1086.
- [171] P.I. Costea, G. Zeller, S. Sunagawa, E. Pelletier, A. Alberti, F. Levenez, M. Tramontano, M. Driessen, R. Hercog, F.E. Jung, J.R. Kultima, M.R. Hayward, L.P. Coelho, E. Allen-Vereece, L. Bertrand, M. Blaut, J.R.M. Brown, T. Carton, S. Cools-Portier, M. Daigneault, M. Derrien, A. Druesne, W.M. de Vos, B.B. Finlay, H.J. Flint, F. Guarner, M. Hattori, H. Heilig, R.A. Luna, J. van Hylckama Vlieg, J. Junick, I. Klymiuk, P. Langella, E. Le Chatelier, V. Mai, C. Manichanh, J.C. Martin, C. Mery, H. Morita, P.W. O'Toole, C. Orvain, K.R. Patil, J. Penders, S. Persson, N. Pons, M. Popova, A. Salonen, D. Saulnier, K.P. Scott, B. Singh, K. Slezak, P. Veiga, J. Versalovic, L. Zhao, E.G. Zoetendal, S.D. Ehrlich, J. Dore, P. Bork, Towards standards for human fecal sample processing in metagenomic studies, *Nat. Biotechnol.* 35 (11) (2017) 1069–1076.
- [172] S.J. Salter, M.J. Cox, E.M. Turek, S.T. Calus, W.O. Cookson, M.F. Moffatt, P. Turner, J. Parkhill, N.J. Loman, A.W. Walker, Reagent and laboratory contamination can critically impact sequence-based microbiome analyses, *BMC Biol.* 12 (2014) 87.
- [173] A. Amir, D. McDonald, J.A. Navas-Molina, J. Debelius, J.T. Morton, E. Hyde, A. Robbins-Pianka, R. Knight, Correcting for microbial blooms in fecal samples during room-temperature shipping, *mSystems* 2 (2) (2017).
- [174] F. Fouhy, J. Deane, M.C. Rea, O. Sullivan, R.P. Ross, G. O'Callaghan, B.J. Plant, C. Stanton, The effects of freezing on faecal microbiota as determined using MiSeq sequencing and culture-based investigations, *PLoS One* 10 (3) (2015), e0119355.
- [175] M. Panek, H. Cipic Paljetak, A. Baresic, M. Peric, M. Matijasic, I. Lojkic, D. Vranesic Bender, Z. Krznaric, D. Verbanac, Methodology challenges in studying human gut microbiota - effects of collection, storage, DNA extraction and next generation sequencing technologies, *Sci. Rep.* 8 (1) (2018) 5143.
- [176] NIH Human Microbiome Portfolio Analysis Team, A review of 10 years of human microbiome research activities at the US National Institutes of Health, *Fiscal Years 2007–2016*, *Microbiome* 7 (1) (2019) 31.
- [177] J.R. Zaneveld, C. Lozupone, J.I. Gordon, R. Knight, Ribosomal RNA diversity predicts genome diversity in gut bacteria and their relatives, *Nucleic Acids Res.* 38 (12) (2010) 3869–3879.
- [178] S. Okuda, Y. Tsuchiya, C. Kiriyama, M. Itoh, H. Morisaki, Virtual metagenome reconstruction from 16S rRNA gene sequences, *Nat. Commun.* 3 (2012) 1203.
- [179] M.G. Langille, J. Zaneveld, J.G. Caporaso, D. McDonald, D. Knights, J.A. Reyes, J.C. Clemente, D.E. Burkepile, R.L. Vega Thurber, R. Knight, R.G. Beiko, C. Huttenhower, Predictive functional profiling of microbial communities using 16S rRNA marker gene sequences, *Nat. Biotechnol.* 31 (9) (2013) 814–821.
- [180] A.W. Walker, J.C. Martin, P. Scott, J. Parkhill, H.J. Flint, K.P. Scott, 16S rRNA gene-based profiling of the human infant gut microbiota is strongly influenced by sample processing and PCR primer choice, *Microbiome* 3 (2015) 26.
- [181] R. Bonnet, A. Suau, J. Dore, G.R. Gibson, M.D. Collins, Differences in rDNA libraries of faecal bacteria derived from 10- and 25-cycle PCRs, *Int. J. Syst. Evol. Microbiol.* 52 (Pt 3) (2002) 757–763.
- [182] W.A. Walters, J.G. Caporaso, C.L. Lauber, D. Berg-Lyons, N. Fierer, R. Knight, PrimerProspector: de novo design and taxonomic analysis of barcoded polymerase chain reaction primers, *Bioinformatics* 27 (8) (2011) 1159–1161.
- [183] L.R. Thompson, J.G. Sanders, D. McDonald, A. Amir, J. Ladau, K.J. Locey, R.J. Prill, A. Tripathi, S.M. Gibbons, G. Ackermann, J.A. Navas-Molina, S. Janssen, E. Kopylova, Y. Vazquez-Baeza, A. Gonzalez, J.T. Morton, S. Mirarab, Z. Zech Xu, L. Jiang, M.F. Haroon, J. Kanbar, Q. Zhu, S. Jin Song, T. Kosciolak, N.A. Bokulich, J. Lefler, C.J. Brislawn, G. Humphrey, S.M. Owens, J. Hampton-Marcell, D. Berg-Lyons, V. McKenzie, N. Fierer, J.A. Fuhrman, A. Clauset, R.L. Stevens, A. Shade, K.S. Pollard, K.D. Goodwin, J.K. Jansson, J.A. Gilbert, R. Knight, C. Earth Microbiome Project, A communal catalogue reveals Earth's multiscale microbial diversity, *Nature* 551 (7681) (2017) 457–463.

- [184] J.M. Janda, S.L. Abbott, 16S rRNA gene sequencing for bacterial identification in the diagnostic laboratory: pluses, perils, and pitfalls, *J. Clin. Microbiol.* 45 (9) (2007) 2761–2764.
- [185] T.A. Nelson, S. Holmes, A.V. Alekseyenko, M. Shenoy, T. Desantis, C.H. Wu, G.L. Andersen, J. Winston, J. Sonnenburg, P.J. Pasricha, A. Spormann, PhyloChip microarray analysis reveals altered gastrointestinal microbial communities in a rat model of colonic hypersensitivity, *Neuro Gastroenterol. Motil.* 23 (2) (2011) 169–177, e141–162.
- [186] F. Godoy-Vitorino, K.C. Goldfarb, E.L. Brodie, M.A. Garcia-Amado, F. Michelangeli, M.G. Dominguez-Bello, Developmental microbial ecology of the crop of the folivorous hoatzin, *ISME J.* 4 (5) (2010) 611–620.
- [187] A.M. Eren, L. Maignain, W.J. Sul, L.G. Murphy, S.L. Grim, H.G. Morrison, M.L. Sogin, Oligotyping: differentiating between closely related microbial taxa using 16S rRNA gene data, *Methods Ecol. Evol.* 4 (12) (2013).
- [188] A. Amir, D. McDonald, J.A. Navas-Molina, E. Kopylova, J.T. Morton, Z. Zech Xu, E.P. Kightley, L.R. Thompson, E.R. Hyde, A. Gonzalez, R. Knight, Deblur rapidly resolves single-nucleotide community sequence patterns, *mSystems* 2 (2) (2017).
- [189] B.J. Callahan, P.J. McMurdie, M.J. Rosen, A.W. Han, A.J. Johnson, S.P. Holmes, DADA2: high-resolution sample inference from Illumina amplicon data, *Nat. Methods* 13 (7) (2016) 581–583.
- [190] K.P. Asshauer, B. Wemheuer, R. Daniel, P. Meinicke, Tax4Fun: predicting functional profiles from metagenomic 16S rRNA data, *Bioinformatics* 31 (17) (2015) 2882–2884.
- [191] S.R. Jun, M.S. Robeson, L.J. Hauser, C.W. Schadt, A.A. Gorin, PanPP: pan-genome-based functional profiles for microbial communities, *BMC Res. Notes* 8 (2015) 479.
- [192] C. Quast, E. Pruesse, P. Yilmaz, J. Gerken, T. Schweer, P. Yarza, J. Peplies, F.O. Glockner, The SILVA ribosomal RNA gene database project: improved data processing and web-based tools, *Nucleic Acids Res.* 41 (Database issue) (2013) D590–D596.
- [193] D. McDonald, M.N. Price, J. Goodrich, E.P. Nawrocki, T.Z. DeSantis, A. Probst, G.L. Andersen, R. Knight, P. Hugenholtz, An improved Greengenes taxonomy with explicit ranks for ecological and evolutionary analyses of bacteria and archaea, *ISME J.* 6 (3) (2012) 610–618.
- [194] T.R. Lesker, A.C. Durairaj, E.J.C. Galvez, I. Lagkouvardos, J.F. Baines, T. Clavel, A. Sczryba, A.C. McHardy, T. Strowig, An integrated metagenome catalog reveals new insights into the murine gut microbiome, *Cell Rep.* 30 (9) (2020) 2909–2922, e2906.
- [195] M. Balvociute, D.H. Huson, R.D.P. Silva, Greengenes, NCBI and OTT – how do these taxonomies compare? *BMC Genom.* 18 (Suppl 2) (2017) 114.
- [196] M. Scholz, D.V. Ward, E. Pasolli, T. Tolio, M. Zolfo, F. Asnicar, D.T. Truong, A. Tett, A.L. Morrow, N. Segata, Strain-level microbial epidemiology and population genomics from shotgun metagenomics, *Nat. Methods* 13 (5) (2016) 435–438.
- [197] S. Mukherjee, R. Seshadri, N.J. Varghese, E.A. Eloe-Fadrosh, J.P. Meier-Kolthoff, M. Goker, R.C. Coates, M. Hadjithomas, G.A. Pavlopoulos, D. Paez-Espino, Y. Yoshikuni, A. Visel, W.B. Whitman, G.M. Garrity, J.A. Eisen, P. Hugenholtz, A. Pati, N.N. Ivanova, T. Woyke, H.P. Klenk, N.C. Kyprides, 1,003 reference genomes of bacterial and archaeal isolates expand coverage of the tree of life, *Nat. Biotechnol.* 35 (7) (2017) 676–683.
- [198] S.G. Tringe, C. von Mering, A. Kobayashi, A.A. Salamov, K. Chen, H.W. Chang, M. Podar, J.M. Short, E.J. Mathur, J.C. Detter, P. Bork, P. Hugenholtz, E.M. Rubin, Comparative metagenomics of microbial communities, *Science* 308 (5721) (2005) 554–557.
- [199] G.W. Tyson, J. Chapman, P. Hugenholtz, E.E. Allen, R.J. Ram, P.M. Richardson, V.V. Solovyev, E.M. Rubin, D.S. Rokhsar, J.F. Banfield, Community structure and metabolism through reconstruction of microbial genomes from the environment, *Nature* 428 (6978) (2004) 37–43.
- [200] C. Quince, A.W. Walker, J.T. Simpson, N.J. Loman, N. Segata, Shotgun metagenomics, from sampling to analysis, *Nat. Biotechnol.* 35 (9) (2017) 833–844.
- [201] F.P. Breitwieser, J. Lu, S.L. Salzberg, A review of methods and databases for metagenomic classification and assembly, *Briefings Bioinf.* 20 (4) (2019) 1125–1136.
- [202] S. Rampelli, M. Candela, S. Turroni, E. Biagi, S. Collino, C. Franceschi, P.W. O'Toole, P. Brigidi, Functional metagenomic profiling of intestinal microbiome in extreme ageing, *Aging (Albany NY)* 5 (12) (2013) 902–912.
- [203] J.L. Round, S.K. Mazmanian, The gut microbiota shapes intestinal immune responses during health and disease, *Nat. Rev. Immunol.* 9 (5) (2009) 313–323.
- [204] Y. Guigoz, J. Dore, E.J. Schiffrin, The inflammatory status of old age can be nurtured from the intestinal environment, *Curr. Opin. Clin. Nutr. Metab. Care* 11 (1) (2008) 13–20.
- [205] E. Pasolli, F. Asnicar, S. Manara, M. Zolfo, N. Karcher, F. Armanini, F. Beghini, P. Manghi, A. Tett, P. Ghensi, M.C. Collado, B.L. Rice, C. DuLong, X.C. Morgan, C.D. Golden, C. Quince, C. Huttenhower, N. Segata, Extensive unexplored human microbiome diversity revealed by over 150,000 genomes from metagenomes spanning age, geography, and lifestyle, *Cell* 176 (3) (2019) 649–662 e620.
- [206] J.B. Emerson, R.I. Adams, C.M.B. Roman, B. Brooks, D.A. Coil, K. Dahlhausen, H.H. Ganz, E.M. Hartmann, T. Hsu, N.B. Justice, I.G. Paulino-Lima, J.C. Luongo, D.S. Lymeropoulou, C. Gomez-Silvan, B. Rothschild-Mancinelli, M. Balk, C. Huttenhower, A. Nocker, P. Vaishampayan, L.J. Rothschild, Schrodinger's microbes: tools for distinguishing the living from the dead in microbial ecosystems, *Microbiome* 5 (1) (2017) 86.
- [207] S. Bashiardes, G. Zilberman-Schapira, E. Elinav, Use of metatranscriptomics in microbiome research, *Bioinf. Biol. Insights* 10 (2016) 19–25.
- [208] M. Shakya, C.C. Lo, P.S.G. Chain, Advances and challenges in metatranscriptomic analysis, *Front. Genet.* 10 (2019) 904.
- [209] G.S. Abu-Ali, R.S. Mehta, J. Lloyd-Price, H. Mallick, T. Branck, K.L. Ivey, D.A. Drew, C. DuLong, E. Rimm, J. Izard, A.T. Chan, C. Huttenhower, Metatranscriptome of human faecal microbial communities in a cohort of adult men, *Nat. Microbiol.* 3 (3) (2018) 356–366.
- [210] J.L. Klassen, Defining microbiome function, *Nat. Microbiol.* 3 (8) (2018) 864–869.
- [211] E.A. Grice, H.H. Kong, G. Renaud, A.C. Young, , N.C.S. Program, G.G. Bouffard, R.W. Blakesley, T.G. Wolfsberg, M.L. Turner, J.A. Segre, A diversity profile of the human skin microbiota, *Genome Res.* 18 (7) (2008) 1043–1050.
- [212] H.J. Kim, J.J. Kim, N.R. Myeong, T. Kim, D. Kim, S. An, H. Kim, T. Park, S.I. Jang, J.H. Yeon, I. Kwack, W.J. Sul, Segregation of age-related skin microbiome characteristics by functionality, *Sci. Rep.* 9 (1) (2019) 16748.
- [213] N. Shibagaki, W. Suda, C. Clavaud, P. Bastien, L. Takayasu, E. Iioka, R. Kurokawa, N. Yamashita, Y. Hattori, C. Shindo, L. Breton, M. Hattori, Aging-related changes in the diversity of women's skin microbiomes associated with oral bacteria, *Sci. Rep.* 7 (1) (2017) 10567.
- [214] L. Gao, T. Xu, G. Huang, S. Jiang, Y. Gu, F. Chen, Oral microbiomes: more and more importance in oral cavity and whole body, *Protein Cell* 9 (5) (2018) 488–500.
- [215] N. Segata, S.K. Haake, P. Mannon, K.P. Lemon, L. Waldron, D. Gevers, C. Huttenhower, J. Izard, Composition of the adult digestive tract bacterial microbiome based on seven mouth surfaces, tonsils, throat and stool samples, *Genome Biol.* 13 (6) (2012) R42.
- [216] F. Galkin, P. Mamoshina, A. Aliper, E. Putin, V. Moskalev, V.N. Gladyshev, A. Zhavoronkov, Human gut microbiome aging clock based on taxonomic profiling and deep learning, *iScience* 23 (6) (2020) 101199.
- [217] P.A. Mackowiak, Recycling metchnikoff: probiotics, the intestinal microbiome and the quest for long life, *Front. Publ. Health* 1 (2013) 52.
- [218] Y. Suzuki, K. Ikeda, K. Sakuma, S. Kawai, K. Sawaki, T. Asahara, T. Takahashi, H. Tsuji, K. Nomoto, R. Nagpal, C. Wang, S. Nagata, Y. Yamashiro, Association between yogurt consumption and intestinal microbiota in healthy young adults differs by host gender, *Front. Microbiol.* 8 (2017) 847.
- [219] D.S. Wiesenborn, E.J.C. Galvez, L. Spinelli, B. Victoria, B. Allen, A. Schneider, A. Gesing, K.A. Al-Regaiey, T. Strowig, K.H. Schafer, M.M. Masternak, The role of ames dwarfism and calorie restriction on gut microbiota, *J. Gerontol. A Biol. Sci. Med. Sci.* 75 (7) (2020) e1–e8.
- [220] C. Fraumene, V. Manghina, E. Cadoni, F. Marongiu, M. Abbondio, M. Serra, A. Palomba, A. Tanca, E. Laconi, S. Uzzau, Caloric restriction promotes rapid expansion and long-lasting increase of Lactobacillus in the rat fecal microbiota, *Gut Microb.* 9 (2) (2018) 104–114.
- [221] A. Tanca, M. Abbondio, A. Palomba, C. Fraumene, F. Marongiu, M. Serra, D. Pagnozzi, E. Laconi, S. Uzzau, Caloric restriction promotes functional changes involving short-chain fatty acid biosynthesis in the rat gut microbiota, *Sci. Rep.* 8 (1) (2018) 14778.
- [222] F. Cabreiro, C. Au, K.Y. Leung, N. Vergara-Irigaray, H.M. Cocheme, T. Noori, D. Weinkove, E. Schuster, N.D. Greene, D. Gems, Metformin retards aging in *C. elegans* by altering microbial folate and methionine metabolism, *Cell* 153 (1) (2013) 228–239.
- [223] A. Zhernakova, A. Kurilshikov, M.J. Bonder, E.F. Tigchelaar, M. Schirmer, T. Vatanen, Z. Mujagic, A.V. Vila, G. Falony, S. Vieira-Silva, J. Wang, F. Imhann, E. Brandsma, S.A. Jankipersadsing, M. Joossens, M.C. Cenit, P. Deelen, M.A. Swertz, s. LifeLines cohort, R.K. Weersma, E.J. Feskens, M.G. Netea, D. Gevers, D. Jonkers, L. Franke, Y.S. Ualchenko, C. Huttenhower, J. Raes, M.H. Hofker, R.J. Xavier, C. Wijmenga, J. Fu, Population-based metagenomics analysis reveals markers for gut microbiome composition and diversity, *Science* 352 (6285) (2016) 565–569.
- [224] K. Forslund, F. Hildebrand, T. Nielsen, G. Falony, E. Le Chatelier, S. Sunagawa, E. Prifti, S. Vieira-Silva, V. Gudmundsdottir, H.K. Pedersen, M. Arumugam, K. Kristiansen, A.Y. Voigt, H. Vestergaard, R. Hercog, P.J. Costea, J.R. Kultima, J. Li, T. Jorgensen, F. Levenez, J. Dore, H.I.T.c. Meta, H.B. Nielsen, S. Brunak, J. Raes, T. Hansen, J. Wang, S.D. Ehrlich, P. Bork, O. Pedersen, Disentangling type 2 diabetes and metformin treatment signatures in the human gut microbiota, *Nature* 528 (7581) (2015) 262–266.
- [225] T. Bryrup, C.W. Thomsen, T. Kern, K.H. Allin, I. Brandslund, N.R. Jorgensen, H. Vestergaard, T. Hansen, T.H. Hansen, O. Pedersen, T. Nielsen, Metformin-induced changes of the gut microbiota in healthy young men: results of a non-blinded, one-armed intervention study, *Diabetologia* 62 (6) (2019) 1024–1035.
- [226] H. Lee, Y. Lee, J. Kim, J. An, S. Lee, H. Kong, Y. Song, C.K. Lee, K. Kim, Modulation of the gut microbiota by metformin improves metabolic profiles in aged obese mice, *Gut Microb.* 9 (2) (2018) 155–165.
- [227] A. Bitto, T.K. Ito, V.V. Pineda, N.J. LeTexier, H.Z. Huang, E. Sutlief, H. Tung, N. Vizzini, B. Chen, K. Smith, Transient Rapamycin Treatment Can Increase Lifespan and Healthspan in Middle-Aged Mice, *elife*, 2016, p. 5, e16351.
- [228] V. Hurez, V. Dao, A. Liu, S. Pandeswara, J. Gelfond, L. Sun, M. Bergman, C.J. Orihuela, V. Galvan, A. Padron, J. Drerup, Y. Liu, P. Hasty, Z.D. Sharp, T.J. Curiel, Chronic mTOR inhibition in mice with rapamycin alters T, B, myeloid, and innate lymphoid cells and gut flora and prolongs life of immune-deficient mice, *Aging Cell* 14 (6) (2015) 945–956.
- [229] M.J. Jung, J. Lee, N.R. Shin, M.S. Kim, D.W. Hyun, J.H. Yun, P.S. Kim,

- T.W. Whon, J.W. Bae, Chronic repression of mTOR complex 2 induces changes in the gut microbiota of diet-induced obese mice, *Sci. Rep.* 6 (2016) 30887.
- [230] T.S. Ghosh, S. Rampelli, I.B. Jeffery, A. Santoro, M. Neto, M. Capri, E. Giampieri, A. Jennings, M. Candela, S. Turroni, E.G. Zoetendal, G.D.A. Hermes, C. Elodie, N. Meunier, C.M. Brugere, E. Pujos-Guillot, A.M. Berendsen, L. De Groot, E.J.M. Feskens, J. Kaluza, B. Pietruszka, M.J. Bielak, B. Comte, M. Maijo-Ferre, C. Nicoletti, W.M. De Vos, S. Fairweather-Tait, A. Cassidy, P. Brigidi, C. Franceschi, P.W. O'Toole, Mediterranean diet intervention alters the gut microbiome in older people reducing frailty and improving health status: the NU-AGE 1-year dietary intervention across five European countries, *Gut* 69 (7) (2020) 1218–1228.
- [231] B.J. Smith, R.A. Miller, A.C. Ericsson, D.C. Harrison, R. Strong, T.M. Schmidt, Changes in the gut microbiome and fermentation products concurrent with enhanced longevity in acarbose-treated mice, *BMC Microbiol.* 19 (1) (2019) 130.